

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Alameda	Alameda County Medical Center	14.0% (8;112) Worse	9.1% (13;159)	22.4% (7;44)	3.6% (3;106)	29.9% (3;9)	2.8% (2;133)	4.0% (5;386)	4.9% (1;23)	2.5% (2;170)
	Alameda Hospital	9.3% (4;24)	9.5% (8;83)	25.4% (4;12)	4.9% (4;71)		3.0% (3;83)	3.5% (5;139)	0.0% (0;30)	4.7% (8;143)
	Alta Bates Summit Medical Center – Alta Bates Campus	4.6% (2;21)	7.6% (25;301)	22.1% (24;104)	0.9% (1;194) Better	0.0% (0;3)	2.6% (4;185)	4.5% (11;268)	0.0% (0;99)	3.5% (6;179)
	Alta Bates Summit Medical Center – Summit Campus – Hawthorne	6.7% (24;418)	8.3% (15;283)	22.8% (6;23)	4.3% (9;260)		2.8% (8;304)	2.0% (11;507)	1.0% (1;100)	5.1% (15;315)
	Eden Medical Center	8.1% (7;43)	9.5% (64;456)	17.4% (24;126)	6.6% (23;281)	26.3% (17;49)	1.7% (3;140)	2.8% (8;235)	2.1% (3;118)	5.2% (10;174)
	Kaiser Foundation Hospital – Hayward	3.8% (8;155)	7.1% (22;264)	16.4% (8;40)	3.8% (11;217)	20.4% (3;7)	2.3% (5;242)	3.5% (15;326)	4.5% (8;148)	4.6% (15;223)
	Kaiser Foundation Hospital – Oakland Campus	4.5% (8;130)	6.6% (29;385) Better	11.3% (9;57) Better	4.7% (19;320)	9.1% (1;8)	2.7% (7;239)	4.0% (24;571)	4.5% (6;152)	5.1% (6;126)
	Saint Rose Hospital	9.8% (18;190) Worse	5.3% (2;63)	14.6% (1;5)	2.6% (1;58)		4.9% (7;132)	4.7% (10;278)	0.0% (0;43)	3.2% (5;169)
	San Leandro Hospital	4.4% (2;24)	4.8% (2;42)	18.3% (1;5)	1.9% (1;37)		0.0% (0;88)	2.6% (4;186)		8.9% (4;64)
	Valleycare Medical Center	7.0% (9;132)	10.8% (4;65)	28.5% (2;5)	5.7% (2;60)		1.0% (1;94)	1.7% (6;234)	0.0% (0;88)	2.7% (7;214)
	Washington Hospital – Fremont	4.5% (9;234)	8.7% (23;261)	20.3% (12;53)	4.9% (8;191)	22.1% (3;17)	4.3% (10;250)	4.5% (17;387)	3.3% (2;85)	5.0% (18;317)
Amador	Sutter Amador Hospital	9.1% (2;13)	11.8% (4;64)	23.8% (1;7)	5.5% (2;55)		4.3% (3;79)	3.6% (3;79)	1.6% (1;53)	2.7% (3;111)
Butte	Biggs Gridley Memorial Hospital		68.2% (1;5)		36.7% (1;5)		0.0% (0;10)	0.0% (0;28)		12.4% (4;87)
	Enloe Medical Center – Esplanade Campus	6.4% (16;334)	10.2% (27;339)	26.9% (17;66)	4.9% (10;272)		1.1% (3;262)	5.5% (19;300) Worse	1.7% (3;151)	5.2% (24;455)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Butte (continued)	Feather River Hospital	5.1% (5;47)	2.1% (1;72)		1.3% (1;70)		2.9% (4;116)	1.8% (4;132)	0.0% (0;75)	0.9% (2;144) Better
	Oroville Hospital	2.9% (2;34)	5.4% (3;95)		3.8% (3;91)		0.7% (1;156)	1.6% (5;262)	1.1% (1;56)	1.4% (2;154)
Calaveras	Mark Twain Medical Center*		13.6% (3;22)		6.8% (2;21)		5.7% (1;22)	1.7% (1;54)	2.9% (1;20)	5.8% (4;58)
Colusa	Colusa Regional Medical Center		66.7% (1;5)		0.0% (0;4)		7.3% (1;11)	5.5% (1;27)		6.0% (2;44)
Contra Costa	Contra Costa Regional Medical Center	6.0% (1;25)	7.0% (1;66)		4.1% (1;66)		1.5% (1;92)	0.9% (1;243)	0.0% (0;11)	2.5% (3;175)
	Doctors Medical Center – San Pablo	5.4% (11;176)	10.4% (7;144)	14.0% (1;5)	6.8% (6;139)		4.0% (5;132)	3.2% (10;441)	0.0% (0;55)	4.9% (8;183)
	John Muir Medical Center – Concord Campus	5.4% (15;342)	13.7% (15;179)	30.4% (5;12)	8.3% (10;166)		2.5% (6;225)	2.8% (11;329)	5.1% (3;61)	3.1% (4;144)
	John Muir Medical Center – Walnut Creek Campus	5.7% (9;97)	10.4% (73;460)	20.6% (34;145)	5.6% (20;273)	39.0% (19;42) Worse	1.5% (5;335)	2.9% (15;357)	5.9% (11;188) Worse	2.8% (6;214)
	Kaiser Foundation Hospital – Antioch	2.3% (3;100)	6.2% (12;175)	14.8% (6;26)	3.7% (6;147)		3.9% (6;139)	3.0% (9;262)	3.2% (3;73)	2.7% (3;100)
	Kaiser Foundation Hospital – Walnut Creek	6.0% (15;224)	5.8% (12;243)	6.2% (2;37) Better	4.1% (9;202)	20.5% (1;4)	0.4% (1;268) Better	3.2% (13;343)	1.1% (2;200)	3.6% (7;188)
	San Ramon Regional Medical Center	10.9% (8;68)	18.1% (11;101) Worse	29.5% (3;6)	12.9% (7;92) Worse	56.2% (1;3)	2.8% (2;96)	4.6% (4;74)	2.7% (1;42)	5.2% (5;129)
	Sutter Delta Medical Center	6.1% (14;173)	2.0% (1;92)	22.5% (1;4)	0.0% (0;88) Better		0.0% (0;135)	3.8% (14;333)	1.8% (1;59)	2.8% (5;194)
Del Norte	Sutter Coast Hospital	12.5% (5;16)	14.7% (9;64)	24.4% (3;6)	10.6% (6;58)		0.0% (0;87)	4.7% (6;82)	0.0% (0;4)	4.4% (9;170)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
El Dorado	Barton Memorial Hospital	0.0% (0;4)	22.7% (1;10)		21.1% (1;9)		1.9% (1;40)	3.2% (1;58)	0.0% (0;24)	6.4% (3;78)
	Marshall Medical Center	10.6% (4;26)	10.0% (8;109)	24.3% (3;9)	5.0% (4;99)		2.1% (3;103)	3.7% (8;145)	0.0% (0;28)	1.4% (3;214)
Fresno	Adventist Medical Center – Reedley	7.2% (1;5)	32.5% (2;13)	83.2% (1;3)	15.6% (1;10)			0.0% (0;20)		11.4% (2;39)
	Clovis Community Medical Center	3.9% (5;115)	9.0% (7;141)	23.8% (1;5)	4.5% (5;135)		2.4% (5;182)	6.2% (14;260) Worse	0.8% (1;132)	3.6% (8;273)
	Coalinga Regional Medical Center		83.9% (1;5)		46.4% (1;5)			6.4% (1;28)		14.2% (1;30)
	Community Regional Medical Center – Fresno	6.6% (31;576)	10.8% (101;775)	26.1% (58;235)	5.4% (25;490)	28.8% (18;50)	1.7% (8;513)	2.4% (15;999)	2.7% (3;113)	4.8% (28;642)
	Fresno Heart and Surgical Hospital	2.9% (1;41)					0.0% (0;28)	2.6% (2;98)		0.0% (0;22)
	Kaiser Foundation Hospital – Fresno	4.6% (6;112)	9.2% (19;199)	21.0% (11;36)	5.7% (8;161)		5.8% (5;106)	3.8% (11;274)	2.9% (4;131)	3.5% (5;134)
	Saint Agnes Medical Center	6.7% (29;418)	11.5% (31;427)	26.1% (9;32)	6.6% (21;392)	24.5% (1;3)	1.7% (10;508)	3.8% (39;884)	2.9% (7;275)	6.0% (41;754) Worse
Glenn	Glenn Medical Center							0.0% (0;18)		11.1% (1;22)
Humboldt	Jerold Phelps Community Hospital									0.0% (0;3)
	Mad River Community Hospital	6.7% (1;7)	17.0% (2;26)		11.7% (2;25)		1.6% (1;51)	0.0% (0;24)	8.6% (1;7)	6.0% (3;66)
	Redwood Memorial Hospital	12.3% (1;6)	9.2% (1;15)	0.0% (0;3)	7.4% (1;12)		0.0% (0;13)	5.6% (2;43)	0.0% (0;19)	0.0% (0;57)
	Saint Joseph Hospital – Eureka	3.5% (8;197)	12.3% (10;105)	35.9% (4;10)	5.6% (5;94)		3.1% (5;114)	3.3% (8;193)	1.7% (2;104)	5.7% (6;117)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Imperial	El Centro Regional Medical Center	7.0% (10;77)	8.8% (7;131)	25.3% (1;5)	4.6% (5;124)		3.2% (6;168)	3.0% (8;250)	2.4% (1;48)	4.1% (10;208)
	Pioneers Memorial Healthcare District	8.1% (3;19)	9.5% (4;75)	39.4% (1;6)	3.2% (2;68)		1.3% (1;92)	2.2% (2;157)	2.8% (1;33)	7.4% (9;150)
Inyo	Northern Inyo Hospital	0.0% (0;4)	0.0% (0;8)		0.0% (0;8)		0.0% (0;17)	0.0% (0;14)	0.0% (0;14)	10.2% (2;29)
	Southern Inyo Hospital									0.0% (0;4)
Kern	Bakersfield Heart Hospital	4.5% (9;245)	0.0% (0;56)		0.0% (0;54)		0.0% (0;40)	2.5% (10;388)		5.3% (8;158)
	Bakersfield Memorial Hospital	5.9% (20;368)	12.1% (25;254)	21.7% (8;46)	7.5% (12;197)	41.7% (5;11)	5.9% (10;198) Worse	5.3% (18;403) Worse	3.9% (4;121)	3.7% (9;287)
	Delano Regional Medical Center	10.1% (2;6)	29.6% (2;12)		32.6% (2;10)		12.4% (4;43) Worse	2.2% (1;67)	0.0% (0;16)	1.8% (3;143)
	Good Samaritan Hospital – Bakersfield						0.0% (0;3)	0.0% (0;35)		5.5% (1;58)
	Kern Medical Center	0.0% (0;5)	12.2% (4;41)	27.6% (2;10)	0.0% (0;27)	45.2% (2;4)	1.8% (1;74)	2.7% (1;102)	4.5% (1;17)	3.1% (1;70)
	Kern Valley Healthcare District	0.0% (0;3)					0.0% (0;9)	0.0% (0;31)		0.0% (0;86)
	Mercy Hospital – Bakersfield	10.1% (6;29)	8.7% (15;184)	22.4% (8;38)	3.7% (5;143)	36.8% (2;3)	2.9% (5;166)	1.8% (3;185)	0.9% (1;145)	4.7% (16;378)
	Ridgecrest Regional Hospital		13.0% (1;14)		0.0% (0;13)		7.0% (1;35)	10.7% (3;47)	0.0% (0;17)	7.7% (4;100)
	San Joaquin Community Hospital	5.3% (17;322)	7.7% (48;544)	19.4% (27;137)	4.1% (17;389)	15.9% (4;18)	0.4% (1;246)	1.5% (9;656) Better	0.8% (1;135)	3.4% (10;406)
	Tehachapi Hospital									0.0% (0;21)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Kings	Adventist Medical Center	7.9% (4;36)	11.0% (7;131)	23.4% (2;8)	6.5% (5;123)		3.3% (6;196)	3.4% (10;363)	5.7% (5;82)	6.8% (22;457) Worse
	Corcoran District Hospital							0.0% (0;7)		0.0% (0;4)
Lake	Saint Helena Hospital – Clearlake		0.0% (0;4)				0.0% (0;18)	0.0% (0;31)		0.0% (0;72)
	Sutter Lakeside Hospital	16.6% (2;8)	0.0% (0;40)		0.0% (0;39)		0.0% (0;28)	0.0% (0;48)	0.0% (0;28)	5.4% (3;87)
Lassen	Banner Lassen Medical Center	11.5% (1;3)	12.5% (1;6)		0.0% (0;5)		0.0% (0;14)	10.6% (1;16)		0.0% (0;44)
Los Angeles	Alhambra Hospital	7.8% (4;28)	4.8% (4;74)	12.4% (2;17)	2.5% (2;57)		1.9% (2;88)	1.5% (2;134)	0.0% (0;29)	2.6% (8;224)
	Antelope Valley Hospital	8.6% (22;345)	12.0% (37;348)	16.6% (10;80)	9.1% (21;255) Worse	34.5% (6;13)	3.0% (8;270)	4.7% (14;390)	6.4% (8;156) Worse	5.7% (17;362)
	Bellflower Medical Center	0.0% (0;4)	0.0% (0;3)		0.0% (0;3)		24.6% (1;7)	0.0% (0;11)		0.0% (0;22)
	Beverly Hospital	5.3% (12;183)	10.3% (15;128)	26.2% (6;19)	4.7% (6;105)	33.7% (3;4)	1.4% (2;155)	2.1% (6;310)	2.0% (1;65)	2.9% (9;232)
	California Hospital Medical Center – Los Angeles	6.4% (5;108)	9.0% (29;267)	19.5% (15;73)	5.1% (10;186)	34.0% (4;8)	2.2% (3;164)	0.4% (1;408) Better	0.0% (0;24)	3.3% (4;203)
	Cedars Sinai Medical Center	8.0% (35;475)	6.4% (52;714) Better	14.3% (20;158) Better	3.3% (18;475)	17.6% (14;81)	2.3% (14;560)	3.5% (46;1,315)	1.6% (4;263)	3.1% (25;732)
	Centinela Hospital Medical Center	4.2% (15;288)	7.2% (22;280)	27.8% (17;57)	1.1% (3;218) Better	37.4% (2;5)	2.2% (9;376)	1.2% (12;1,007) Better	1.5% (1;49)	1.7% (6;326) Better
	Citrus Valley Medical Center – Inter Community Campus	7.9% (23;416)	4.1% (5;113) Better	7.8% (2;26)	0.0% (0;80)	23.3% (3;7)	4.2% (6;148)	4.5% (10;292)	0.0% (0;81)	3.9% (7;176)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Los Angeles (continued)	Citrus Valley Medical Center – Queen of the Valley Campus	5.5% (6;64)	6.6% (13;263)	14.2% (6;47)	3.4% (5;213)	34.0% (2;3)	2.0% (4;203)	1.6% (4;318)	1.8% (1;75)	5.8% (10;224)
	City of Hope Helford Clinical Research Hospital	14.3% (1;6)	13.1% (1;7)		0.0% (0;4)		0.0% (0;24)	28.2% (3;12) Worse		6.7% (7;138)
	Coast Plaza Hospital*	3.5% (3;39)	12.2% (2;17)	29.6% (1;3)	6.6% (1;14)		1.5% (1;56)	2.8% (2;87)	0.0% (0;13)	4.4% (4;100)
	Community Hospital of Long Beach	11.5% (6;39)	0.0% (0;27)		0.0% (0;25)		4.1% (2;60)	6.4% (4;82)	0.0% (0;28)	6.1% (9;132)
	Community and Mission Hospital of Huntington Park – Slauson	0.0% (0;49)	3.7% (1;23)		4.2% (1;21)		7.7% (5;81) Worse	2.1% (3;201)	6.4% (1;11)	8.2% (7;99)
	Downey Regional Medical Center	4.0% (11;277)	5.8% (7;165)	19.1% (4;28)	2.1% (2;125)	14.0% (1;12)	1.6% (4;249)	2.6% (10;391)	1.7% (2;110)	2.0% (6;286)
	East Los Angeles Doctors Hospital	8.0% (4;36)	11.2% (1;16)		7.3% (1;15)		5.4% (5;46)	0.0% (0;87)	0.0% (0;11)	7.7% (8;87)
	East Valley Hospital Medical Center	0.0% (0;13)	0.0% (0;11)		0.0% (0;11)		6.6% (1;15)	0.0% (0;16)	25.3% (1;7)	3.3% (6;135)
	Encino Hospital Medical Center	3.4% (2;31)	5.2% (2;18)		1.7% (1;17)		2.3% (2;42)	0.4% (1;100) Better	0.0% (0;22)	1.6% (1;41)
	Foothill Presbyterian Hospital – Johnston Memorial	6.1% (3;28)	4.9% (3;72)	9.7% (1;12)	3.0% (2;59)		5.1% (5;106)	4.6% (6;154)	1.5% (1;64)	2.6% (3;139)
	Garfield Medical Center	6.2% (14;227)	6.0% (18;301) Better	13.4% (10;81)	3.4% (6;211)	24.4% (2;9)	2.8% (4;186)	2.7% (12;426)	2.7% (2;71)	4.2% (20;357)
	Glendale Adventist Medical Center – Wilson Terrace	3.3% (12;354) Better	6.8% (25;336)	21.8% (15;65)	2.0% (5;245) Better	18.1% (5;26)	1.2% (4;287)	1.8% (11;471)	0.0% (0;118)	3.0% (23;545)
	Glendale Memorial Hospital and Medical Center	8.1% (20;224)	3.4% (3;96) Better	8.3% (2;21)	1.8% (1;75)		0.8% (1;139)	2.6% (12;468)	3.2% (3;84)	3.5% (13;323)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Los Angeles (continued)	Good Samaritan Hospital – Los Angeles	7.6% (25;315)	7.6% (8;148)	27.2% (5;31)	2.9% (3;114)	0.0% (0;3)	3.7% (6;167)	3.0% (12;405)	0.0% (0;68)	5.9% (18;236)
	Greater El Monte Community Hospital	12.2% (3;14)	8.6% (1;19)		0.0% (0;18)		0.0% (0;44)	3.6% (2;71)		3.2% (6;126)
	Henry Mayo Newhall Memorial Hospital	3.9% (8;203)	9.2% (13;211)	21.1% (5;28)	5.6% (7;174)	19.5% (1;9)	1.9% (5;238)	1.8% (7;320)	5.8% (4;75)	2.0% (8;340)
	Hollywood Presbyterian Medical Center	7.7% (20;218)	10.0% (31;216)	20.2% (16;63)	7.1% (12;145)	27.8% (3;8)	2.0% (5;181)	4.4% (9;286)	0.9% (1;78)	4.1% (8;173)
	Huntington Memorial Hospital	6.4% (21;272)	8.9% (44;426)	20.1% (15;72)	4.0% (14;311)	30.0% (15;43)	1.5% (4;295)	3.2% (14;451)	1.0% (2;223)	3.2% (14;392)
	Kaiser Foundation Hospital – Baldwin Park	3.7% (4;97)	7.9% (14;310)	19.5% (4;26)	4.0% (9;282)		1.5% (3;267)	2.6% (9;503)	1.0% (1;121)	2.3% (7;321)
	Kaiser Foundation Hospital – Downey	1.6% (2;123) Better	6.2% (10;241)	11.7% (4;48)	3.2% (4;191)		2.3% (5;325)	2.5% (9;420)	1.3% (1;104)	4.3% (15;404)
	Kaiser Foundation Hospital – Panorama City	4.9% (6;96)	11.7% (14;201)	18.2% (2;19)	7.2% (12;181)		3.8% (8;203)	5.2% (16;286) Worse	0.0% (0;133)	5.2% (21;346)
	Kaiser Foundation Hospital – South Bay	5.1% (6;86)	8.6% (12;210)	9.5% (2;21)	6.1% (10;189)		3.6% (8;258)	3.0% (10;374)	4.8% (1;33)	3.8% (7;216)
	Kaiser Foundation Hospital – Sunset	3.8% (30;1,125) Better	7.4% (54;669)	12.6% (23;270) Better	6.5% (20;298)	20.1% (11;101)	1.9% (4;197)	2.8% (15;600)	1.7% (1;68)	3.1% (3;96)
	Kaiser Foundation Hospital – West Los Angeles	3.7% (5;108)	6.6% (13;305)	14.9% (3;20)	3.8% (10;284)		2.3% (8;315)	1.6% (5;329)	2.3% (1;44)	3.3% (10;198)
	Kaiser Foundation Hospital – Woodland Hills	3.0% (4;103)	11.2% (13;192)	25.0% (2;14)	6.7% (11;177)		2.7% (6;186)	5.4% (28;341) Worse	1.2% (3;184)	2.4% (7;221)
	Keck Hospital of University of Southern California	12.3% (13;74) Worse	9.5% (36;254)	24.6% (14;65)	6.5% (11;107)	16.3% (11;82)	1.4% (1;49)	8.2% (17;163) Worse	4.7% (1;13)	2.2% (1;47)
	Lakewood Regional Medical Center	7.4% (24;294)	10.2% (36;315)	25.4% (24;79)	5.0% (11;231)	27.1% (1;5)	4.8% (7;169)	2.6% (7;347)	4.2% (3;99)	4.9% (10;205)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Los Angeles (continued)	Long Beach Memorial Medical Center	8.1% (18;284)	9.0% (43;536)	18.7% (18;106)	4.5% (14;390)	33.7% (11;40)	3.5% (9;303)	4.5% (18;544)	0.0% (0;133)	3.6% (13;447)
	Los Angeles Community Hospital	6.1% (3;44)	0.0% (0;43)		0.0% (0;41)		2.4% (3;109)	1.9% (3;174)	0.0% (0;11)	2.1% (3;151)
	Los Angeles County/Harbor – UCLA Medical Center	8.1% (24;326)	9.4% (21;242)	19.0% (9;54)	4.8% (6;168)	39.2% (6;20)	1.0% (2;205)	2.4% (7;531)	0.0% (0;29)	2.0% (2;160)
	Los Angeles County/Olive View – UCLA Medical Center	5.7% (2;105)	0.0% (0;100) Better	0.0% (0;8)	0.0% (0;91)		3.2% (4;145)	3.3% (5;308)		1.2% (1;183)
	Los Angeles County/University of Southern California Medical Center	8.2% (16;273)	14.7% (70;360) Worse	30.0% (32;123) Worse	11.7% (18;170) Worse	33.0% (20;67)	2.1% (6;324)	5.7% (18;685) Worse	4.0% (3;47)	5.9% (16;503)
	Los Angeles Metropolitan Medical Center							0.0% (0;21)		0.0% (0;15)
	Marina Del Rey Hospital	4.0% (1;18)	11.4% (2;37)		6.2% (2;37)		2.7% (4;93)	1.4% (2;129)	1.9% (1;45)	3.2% (4;120)
	Memorial Hospital of Gardena	4.7% (6;59)	5.0% (1;43)		0.0% (0;42)		2.9% (6;145)	1.4% (2;220)	3.1% (1;20)	2.3% (4;164)
	Methodist Hospital of Southern California	5.6% (20;250)	7.6% (28;359)	22.1% (16;64)	3.4% (11;290)	15.0% (1;5)	2.7% (10;286)	2.0% (11;404)	2.9% (5;172)	3.9% (16;319)
	Mission Community Hospital – Panorama Campus	0.0% (0;31)	4.5% (2;45)	22.3% (2;8)	0.0% (0;35)		0.9% (1;78)	0.9% (2;172)	0.0% (0;28)	3.4% (8;145)
	Monterey Park Hospital	0.0% (0;16)	13.5% (4;46)	37.5% (3;12)	6.0% (1;34)		4.8% (2;66)	2.2% (2;108)	0.0% (0;11)	1.2% (1;94)
	Motion Picture and Television Hospital		11.2% (1;3)							9.0% (1;17)
	Northridge Hospital Medical Center	4.6% (16;283)	7.7% (17;258)	22.0% (8;47)	3.5% (8;208)	32.8% (1;3)	1.4% (3;191)	3.3% (17;392)	0.8% (1;116)	3.1% (10;270)
	Norwalk Community Hospital	4.9% (3;45)	0.0% (0;17)		0.0% (0;16)		0.0% (0;69)	0.0% (0;53)	0.0% (0;12)	0.7% (1;86) Better

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Los Angeles (continued)	Olympia Medical Center	0.0% (0;33)	0.0% (0;48) Better	0.0% (0;4)	0.0% (0;43)		3.3% (4;114)	3.9% (5;144)	3.0% (1;28)	2.3% (4;145)
	Pacific Alliance Medical Center, Inc.	0.0% (0;13)	0.0% (0;86) Better	0.0% (0;6)	0.0% (0;78)		0.9% (1;98)	0.9% (1;165)	0.0% (0;21)	1.6% (1;125)
	Pacific Hospital of Long Beach	6.9% (1;14)	6.5% (1;18)		3.6% (1;18)		2.9% (1;30)	0.0% (0;62)	0.0% (0;15)	0.0% (0;53)
	Pacifica Hospital of the Valley	8.6% (3;13)	18.1% (4;24)	36.5% (2;3)	14.4% (2;20)		9.6% (5;51) Worse	0.0% (0;83)	0.0% (0;15)	2.5% (3;93)
	Palmdale Regional Medical Center	8.0% (11;238)	7.3% (7;115)	8.6% (1;16)	4.5% (3;91)	22.7% (3;8)	4.9% (9;189) Worse	2.8% (10;432)	4.6% (3;76)	5.7% (15;298)
	Pomona Valley Hospital Medical Center	5.1% (17;347)	10.2% (45;406)	27.5% (26;88)	4.4% (14;309)	32.5% (5;9)	2.3% (7;278)	1.5% (8;520) Better	0.0% (0;123)	2.9% (9;286)
	Presbyterian Intercommunity Hospital	5.2% (38;584)	6.1% (32;443) Better	16.5% (16;87)	2.7% (12;342) Better	17.4% (4;14)	2.1% (7;334)	1.2% (7;532) Better	2.3% (5;200)	2.3% (9;364)
	Providence Holy Cross Medical Center	6.4% (17;257)	6.6% (19;283)	23.8% (15;55)	1.7% (4;221) Better	0.0% (0;7)	1.4% (5;302)	2.2% (15;495)	3.2% (4;127)	1.8% (11;404) Better
	Providence Little Company of Mary Medical Center – San Pedro	5.8% (2;19)	8.7% (11;105)	22.4% (6;18)	1.8% (1;79)	32.5% (4;8)	1.9% (2;128)	3.6% (5;164)	0.0% (0;47)	1.6% (2;136)
	Providence Little Company of Mary Medical Center – Torrance	7.1% (33;391)	10.5% (53;414)	27.0% (32;106)	4.6% (17;300)	43.6% (4;8)	2.3% (8;311)	2.8% (18;539)	2.1% (3;162)	5.3% (25;375)
	Providence Saint Joseph Medical Center	6.8% (14;274)	8.6% (39;427)	24.9% (19;84)	3.7% (15;315)	26.8% (5;28)	1.3% (4;336)	2.6% (16;457)	1.3% (2;190)	2.9% (14;411)
	Providence Tarzana Medical Center	5.0% (9;162)	6.9% (7;142)	26.1% (5;23)	2.0% (2;118)		0.8% (2;183)	1.5% (9;379) Better	1.8% (3;126)	4.1% (20;322)
	Ronald Reagan UCLA Medical Center	6.9% (15;218)	8.0% (60;414)	20.3% (30;117)	4.1% (18;229)	17.8% (12;68)	3.1% (6;164)	4.5% (16;347)	2.1% (1;26)	4.2% (8;186)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Los Angeles (continued)	Saint Francis Medical Center	3.1% (4;136)	9.2% (23;198)	26.2% (15;47)	2.4% (3;137)	28.2% (5;14)	2.8% (5;180)	1.6% (6;521)	5.4% (2;45)	2.9% (6;214)
	Saint John's Health Center	6.5% (10;147)	9.5% (11;121)	26.3% (6;28)	3.4% (3;88)	41.7% (2;5)	3.2% (5;179)	4.3% (12;199)	3.1% (5;151)	3.7% (11;271)
	Saint Mary Medical Center	5.2% (11;178)	9.8% (17;152)	24.0% (9;35)	5.0% (7;114)	32.8% (1;3)	0.9% (1;136)	0.9% (2;304)	0.0% (0;50)	3.1% (5;189)
	Saint Vincent Medical Center	5.4% (11;175)	7.7% (5;115)	32.8% (4;20)	1.7% (1;94)		4.3% (5;151)	2.6% (8;340)	0.0% (0;45)	3.0% (7;225)
	San Dimas Community Hospital	6.8% (1;10)	3.2% (1;35)	11.2% (1;6)	0.0% (0;29)		0.0% (0;58)	2.4% (2;89)	0.0% (0;36)	2.9% (2;80)
	San Gabriel Valley Medical Center	6.3% (5;47)	6.0% (4;89)	6.4% (1;23)	6.3% (3;65)		0.6% (1;141)	3.1% (11;250)	1.0% (1;73)	7.0% (20;229) Worse
	Santa Monica – UCLA Medical Center and Orthopedic Hospital	4.6% (7;160)	7.1% (4;89)	0.0% (0;5)	4.4% (4;84)		2.6% (5;211)	2.7% (9;307)	0.5% (1;161)	4.2% (14;312)
	Sherman Oaks Hospital	0.5% (1;116) Better	0.0% (0;21)		0.0% (0;19)		0.5% (1;105)	0.8% (2;134) Better	0.0% (0;32)	1.9% (3;80)
	Silver Lake Medical Center – Downtown Campus	0.0% (0;5)	0.0% (0;14)		0.0% (0;14)		1.8% (1;47)	2.7% (1;53)	0.0% (0;4)	7.7% (6;84)
	Southern California Hospital at Culver City*	3.0% (2;60)	14.8% (6;61)	29.9% (1;3)	8.4% (5;58)		2.7% (5;152)	1.5% (3;236)	0.0% (0;33)	2.9% (8;220)
	Southern California Hospital at Hollywood*	0.0% (0;38) Better	0.0% (0;15)		0.0% (0;15)		1.0% (1;59)	1.5% (1;75)	0.0% (0;4)	4.8% (6;103)
	Temple Community Hospital	0.0% (0;12)	0.0% (0;29)		0.0% (0;28)		0.0% (0;34)	1.8% (1;64)	0.0% (0;5)	2.5% (2;58)
	Torrance Memorial Medical Center	4.0% (20;433) Better	9.4% (37;428)	22.1% (19;79)	5.3% (16;340)	27.7% (2;9)	1.5% (7;451)	2.2% (15;597)	3.0% (7;239)	4.2% (25;477)
	Tri-City Regional Medical Center – Hawaiian Gardens	9.4% (3;22)	7.0% (1;14)	0.0% (0;3)	4.8% (1;11)		5.3% (3;51)	5.5% (4;106)	15.9% (1;7)	5.9% (5;92)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Los Angeles (continued)	University of Southern California Verdugo Hills Hospital*	2.4% (1;33)	4.6% (2;77)	15.9% (1;11)	1.9% (1;66)		2.7% (2;104)	4.6% (5;120)	1.6% (1;68)	4.3% (6;166)
	Valley Presbyterian Hospital	4.8% (12;205)	12.0% (19;174)	26.8% (6;32)	6.6% (12;140)		3.3% (5;148)	2.1% (9;372)	3.3% (2;49)	4.6% (11;184)
	West Hills Hospital and Medical Center	8.3% (16;193)	15.4% (23;164) Worse	6.7% (1;18)	10.9% (20;141) Worse	40.7% (2;5)	5.0% (12;195) Worse	3.6% (12;238)	3.0% (3;117)	6.1% (20;198)
	White Memorial Medical Center	7.5% (23;276)	8.1% (27;282)	19.9% (14;68)	4.5% (12;211)	17.3% (1;3)	1.0% (3;245)	1.6% (5;339)	2.4% (2;62)	3.5% (8;267)
	Whittier Hospital Medical Center	4.6% (4;41)	3.2% (1;64)	0.0% (0;6)	2.3% (1;58)		0.7% (1;133)	2.3% (4;190)	0.0% (0;39)	4.0% (4;160)
Madera	Madera Community Hospital	12.6% (7;47)	36.0% (5;34) Worse	60.4% (1;3)	21.6% (3;30)		0.0% (0;54)	8.5% (7;119) Worse	0.0% (0;10)	13.6% (16;139) Worse
Marin	Kaiser Foundation Hospital – San Rafael	9.2% (7;64)	9.5% (13;156)	36.3% (7;21)	3.2% (5;134)		4.2% (3;108)	2.5% (5;164)	1.6% (2;109)	4.7% (4;64)
	Marin General Hospital	4.0% (6;204)	15.3% (24;181) Worse	36.0% (12;32)	8.5% (12;148)		2.7% (5;159)	4.9% (13;199)	0.0% (0;100)	5.2% (12;221)
	Novato Community Hospital	0.0% (0;5)	10.8% (2;33)		6.3% (2;32)		0.0% (0;38)	1.0% (1;77)	0.0% (0;43)	2.8% (2;67)
Mariposa	John C. Fremont Healthcare District	14.2% (1;6)					0.0% (0;5)	0.0% (0;7)		0.0% (0;20)
Mendocino	Frank R. Howard Memorial Hospital		7.5% (1;25)		0.0% (0;23)		15.0% (1;13)	7.3% (3;42)	2.0% (1;53)	4.5% (1;38)
	Mendocino Coast District Hospital	0.0% (0;3)	10.2% (1;13)		5.4% (1;13)		8.9% (2;29)	5.2% (1;32)	7.0% (1;18)	4.2% (2;57)
	Ukiah Valley Medical Center	0.0% (0;8)	15.2% (7;68)	0.0% (0;3)	9.3% (7;65)		0.0% (0;85)	1.7% (3;132)	8.1% (2;28)	3.0% (5;136)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Merced	Memorial Hospital Los Banos		16.3% (2;5)		11.1% (1;4)			0.0% (0;40)		0.0% (0;35)
	Mercy Medical Center – Merced	4.1% (5;60)	9.9% (13;159)	29.9% (4;9)	4.8% (8;149)		2.9% (5;195)	2.1% (9;351)	2.1% (2;100)	2.8% (14;439)
Modoc	Modoc Medical Center		0.0% (0;4)		0.0% (0;4)			38.3% (1;8)		0.0% (0;29)
Mono	Mammoth Hospital						0.0% (0;5)	0.0% (0;5)	0.0% (0;3)	0.0% (0;8)
Monterey	Community Hospital of The Monterey Peninsula*	9.1% (10;135)	11.7% (22;245)	17.0% (6;37)	8.9% (14;204) Worse	27.5% (2;4)	2.8% (6;221)	4.8% (11;256)	2.7% (4;153)	3.4% (9;287)
	George L. Mee Memorial Hospital							0.0% (0;22)	0.0% (0;5)	0.0% (0;29)
	Natividad Medical Center	0.0% (0;8)	6.9% (1;35)	23.7% (1;5)	0.0% (0;30)		2.3% (1;68)	1.8% (1;98)	14.4% (2;14)	4.5% (5;113)
	Salinas Valley Memorial Hospital	8.6% (21;242)	10.2% (23;232)	23.6% (13;47)	5.9% (10;184)		1.4% (3;193)	2.9% (12;329)	1.9% (2;91)	2.9% (7;188)
Napa	Queen of the Valley Hospital – Napa	8.0% (9;123)	18.1% (23;133) Worse	31.6% (10;33)	15.3% (13;98) Worse		2.3% (3;126)	6.9% (18;263) Worse	2.0% (2;83)	8.6% (13;164) Worse
	Saint Helena Hospital	2.9% (4;141)	3.3% (1;33)		1.9% (1;32)		2.1% (2;70)	0.6% (1;137)	0.0% (0;34)	4.0% (4;80)
Nevada	Sierra Nevada Memorial Hospital	7.1% (6;37)	11.5% (7;80)	38.2% (3;6)	4.1% (3;73)		2.9% (3;133)	3.0% (6;149)	5.3% (6;100) Worse	2.9% (8;244)
	Tahoe Forest Hospital	0.0% (0;5)	0.0% (0;4)		0.0% (0;4)		6.9% (1;20)	0.0% (0;18)	0.0% (0;24)	0.0% (0;39)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Orange	AHMC Anaheim Regional Medical Center	4.7% (15;288)	7.0% (7;163)	17.9% (4;27)	2.9% (2;135)		3.6% (8;243)	3.8% (16;501)	2.9% (2;90)	1.5% (4;321) Better
	Anaheim General Hospital		0.0% (0;3)		0.0% (0;3)		0.0% (0;6)	0.0% (0;6)		9.3% (2;25)
	Chapman Medical Center		11.8% (3;15)	32.3% (3;6)	0.0% (0;9)		0.0% (0;25)	8.6% (2;31)	0.0% (0;10)	6.2% (2;51)
	Coastal Communities Hospital	8.2% (1;10)	24.0% (1;15)		16.5% (1;14)		2.1% (1;39)	8.8% (5;59)	0.0% (0;9)	4.4% (3;82)
	Fountain Valley Regional Hospital and Medical Center – Euclid	5.2% (18;307)	7.5% (30;345)	19.9% (17;75)	3.2% (10;266)	31.0% (3;4)	1.7% (4;253)	1.6% (6;327)	1.5% (1;100)	4.6% (21;426)
	Garden Grove Hospital and Medical Center	11.1% (8;33)	6.5% (4;51)	22.0% (4;16)	0.0% (0;35)		2.1% (3;107)	1.4% (2;103)	0.0% (0;28)	2.6% (5;132)
	Hoag Memorial Hospital Presbyterian	6.1% (24;399)	7.2% (45;602)	16.9% (19;116)	3.8% (19;459)	21.3% (7;27)	1.2% (7;512)	3.7% (16;315)	0.5% (1;140)	1.1% (4;339) Better
	Hoag Orthopedic Institute								0.0% (0;212)	
	Huntington Beach Hospital	0.0% (0;24)	3.5% (1;29)		0.0% (0;28)		1.5% (1;41)	2.1% (3;97)	4.0% (1;14)	5.1% (5;68)
	Kaiser Foundation Hospital – Orange County – Anaheim	3.5% (7;173)	8.5% (37;511)	16.4% (16;135)	5.4% (15;350)	26.8% (6;26)	1.0% (3;390)	3.6% (17;514)	3.5% (8;210)	2.5% (11;477)
	La Palma Intercommunity Hospital	2.6% (2;45)	9.9% (3;48)	0.0% (0;3)	5.5% (2;44)		1.5% (1;57)	2.9% (4;116)	0.0% (0;27)	0.0% (0;100) Better
	Los Alamitos Medical Center	5.4% (9;174)	10.6% (24;254)	28.8% (14;49)	5.0% (10;204)		1.4% (2;169)	4.8% (13;311)	1.9% (3;171)	5.2% (14;302)
	Mission Hospital Laguna Beach	0.0% (0;7)	0.0% (0;3)		0.0% (0;3)		0.0% (0;53)	1.9% (1;48)	0.0% (0;33)	1.8% (1;65)
	Mission Hospital Regional Medical Center	4.5% (14;289)	9.4% (42;378)	20.5% (18;87)	5.0% (14;256)	29.0% (10;35)	3.7% (9;215)	3.4% (14;317)	1.2% (2;163)	4.5% (14;305)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Orange (continued)	Orange Coast Memorial Medical Center	3.9% (8;169)	5.4% (5;136)	15.6% (2;27)	2.0% (2;106)	28.2% (1;3)	2.1% (4;208)	3.0% (11;369)	0.0% (0;104)	6.0% (17;229)
	Placentia Linda Hospital	5.7% (4;35)	9.6% (3;55)	79.0% (1;3)	2.5% (1;51)		1.9% (1;72)	1.8% (2;130)	2.1% (1;58)	4.2% (5;164)
	Saddleback Memorial Medical Center	4.6% (13;270)	11.0% (38;327)	20.2% (12;52)	7.0% (25;273)		1.1% (3;242)	2.5% (20;494)	0.7% (2;248)	2.9% (16;407)
	Saint Joseph Hospital – Orange	5.1% (17;299)	9.2% (32;363)	24.5% (19;79)	5.0% (12;273)	9.0% (1;11)	2.6% (9;289)	3.7% (20;459)	1.0% (2;194)	3.2% (12;336)
	Saint Jude Medical Center	5.0% (21;329)	6.9% (38;453) Better	11.5% (10;92) Better	4.2% (20;331)	25.0% (8;30)	1.6% (4;241)	1.9% (10;394)	1.2% (2;213)	4.6% (17;353)
	UC Irvine Medical Center	12.7% (19;148) Worse	9.8% (73;446)	26.2% (37;120)	4.9% (19;244)	20.0% (17;82)	2.2% (6;215)	4.0% (8;268)	7.2% (4;51)	3.1% (3;98)
	West Anaheim Medical Center	6.8% (11;136)	11.2% (6;55)	21.2% (2;8)	7.2% (4;47)		0.8% (1;97)	2.4% (6;235)	0.0% (0;49)	1.5% (5;246) Better
	Western Medical Center – Anaheim	2.7% (1;79)	10.8% (2;12)		12.0% (1;9)		5.2% (2;37)	10.6% (2;30)	0.0% (0;7)	5.9% (2;45)
	Western Medical Center – Santa Ana	5.8% (10;219)	12.6% (42;222) Worse	23.6% (17;71)	8.9% (15;132) Worse	32.7% (10;19)	4.3% (2;63)	5.6% (4;108)	0.0% (0;40)	2.9% (2;89)
Placer	Kaiser Foundation Hospital – Roseville	6.1% (17;272)	6.6% (14;308)	28.4% (4;14)	3.1% (10;293)		3.4% (8;251)	3.9% (19;397)	3.9% (10;233)	4.2% (7;120)
	Sutter Auburn Faith Hospital	1.7% (1;30)	9.2% (5;96)	36.8% (4;9)	2.0% (1;87)		3.3% (2;80)	2.1% (2;89)	0.8% (1;81)	4.8% (6;129)
	Sutter Roseville Medical Center	6.0% (19;302)	13.4% (55;377) Worse	30.5% (25;77)	8.5% (26;282) Worse	25.0% (4;18)	2.1% (6;335)	3.0% (10;294)	3.2% (6;179)	3.0% (10;356)
Plumas	Eastern Plumas Hospital – Portola Campus	0.0% (0;3)						0.0% (0;3)		11.8% (1;27)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Plumas (continued)	Plumas District Hospital		0.0% (0;3)		0.0% (0;3)		9.1% (1;8)	9.1% (1;11)		0.0% (0;22)
	Seneca Healthcare District		0.0% (0;3)		0.0% (0;3)			42.4% (4;15) Worse		0.0% (0;25)
Riverside	Corona Regional Medical Center – Main	5.9% (3;31)	2.2% (1;83)	33.0% (1;6)	0.0% (0;77) Better		1.5% (3;144)	2.4% (9;280)	2.5% (3;89)	1.9% (7;302) Better
	Desert Regional Medical Center	8.5% (24;424)	8.7% (51;525)	22.8% (31;134)	3.3% (11;339)	29.0% (9;52)	3.2% (8;311)	5.1% (14;412)	0.7% (1;199)	4.3% (12;392)
	Eisenhower Medical Center	5.5% (21;416)	5.0% (20;570) Better	15.1% (10;66)	2.1% (8;491) Better	16.5% (2;13)	0.8% (3;437)	2.1% (13;612)	1.5% (4;278)	3.9% (28;734)
	Hemet Valley Medical Center	4.9% (14;150)	8.9% (15;188)	16.9% (3;10)	5.2% (12;178)		1.8% (5;248)	3.3% (15;428)	2.5% (4;173)	3.3% (16;383)
	John F. Kennedy Memorial Hospital	7.5% (11;127)	3.4% (1;32)		0.0% (0;31)		1.3% (1;71)	5.7% (6;155)	1.6% (1;85)	3.0% (6;213)
	Kaiser Foundation Hospital – Moreno Valley	9.2% (7;41)	6.2% (5;81)	47.0% (2;4)	1.9% (2;75)		1.2% (1;86)	1.6% (2;140)	3.6% (1;27)	3.3% (5;139)
	Kaiser Foundation Hospital – Riverside	9.5% (16;115)	7.5% (7;151)	13.0% (2;23)	4.2% (4;127)		2.9% (4;166)	3.1% (8;284)	4.1% (4;79)	1.7% (3;217)
	Loma Linda University Medical Center – Murrieta	9.0% (12;251)	8.9% (4;72)	32.8% (3;17)	2.6% (1;53)		0.9% (1;129)	2.7% (6;330)	0.0% (0;43)	3.3% (5;196)
	Menifee Valley Medical Center	7.7% (10;58)	5.3% (1;47)		3.1% (1;46)		2.9% (2;69)	0.8% (1;154)	0.0% (0;63)	3.6% (3;111)
	Palo Verde Hospital	19.9% (1;6)	11.4% (1;4)		0.0% (0;3)		0.0% (0;28)	11.2% (1;19)		2.3% (1;55)
	Parkview Community Hospital Medical Center	16.2% (6;33) Worse	9.0% (6;111)		4.2% (4;108)		1.5% (2;147)	3.5% (7;235)	0.0% (0;55)	5.1% (14;273)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Riverside (continued)	Riverside Community Hospital	8.8% (61;837) Worse	8.8% (37;406)	22.2% (19;93)	4.2% (12;298)	26.0% (6;15)	2.9% (9;348)	2.5% (13;546)	2.5% (4;161)	4.1% (22;522)
	Riverside County Regional Medical Center	9.5% (4;51)	10.3% (13;204)	24.5% (8;38)	4.9% (4;162)	99.1% (1;4)	5.6% (6;140)	3.8% (8;402)	5.9% (3;53)	1.4% (2;198)
	San Geronio Memorial Hospital	4.3% (6;107)	12.3% (4;52)	20.1% (1;4)	8.2% (3;48)		2.0% (1;62)	2.1% (3;173)	0.0% (0;21)	4.2% (9;212)
	Southwest Healthcare System – Murrieta	7.3% (11;92)	9.1% (21;284)	24.2% (11;54)	2.9% (5;221)	39.5% (5;9)	2.9% (6;217)	2.9% (10;327)	0.8% (1;181)	2.9% (11;370)
	Temecula Valley Hospital		11.7% (1;9)		0.0% (0;7)		13.5% (1;11)	0.0% (0;11)	0.0% (0;4)	0.0% (0;11)
Sacramento	Kaiser Foundation Hospital – Sacramento	5.1% (9;120)	7.7% (69;539)	15.8% (35;196) Better	7.2% (25;266) Worse	11.7% (9;77) Better	2.5% (5;210)	5.5% (16;244) Worse	2.2% (3;147)	7.1% (6;70)
	Kaiser Foundation Hospital – South Sacramento	2.2% (5;181) Better	9.0% (22;304)	30.0% (13;33)	3.1% (7;268)	31.5% (2;3)	3.6% (5;133)	2.6% (13;434)	0.7% (1;119)	5.0% (7;119)
	Mercy General Hospital	5.0% (43;1,202)	5.0% (16;349) Better	9.4% (5;57) Better	2.6% (7;269)	22.4% (4;23)	3.2% (7;189)	2.2% (13;539)	3.2% (3;97)	4.2% (7;180)
	Mercy Hospital – Folsom	3.5% (1;12)	10.9% (5;76)	34.4% (4;9)	3.1% (1;67)		2.1% (3;139)	1.8% (4;142)	3.1% (3;63)	1.7% (2;93)
	Mercy San Juan Hospital	6.8% (28;318)	9.6% (109;711)	22.6% (55;203)	5.4% (41;454)	22.1% (13;54)	2.0% (7;316)	2.3% (20;643)	1.9% (4;179)	2.5% (8;321)
	Methodist Hospital of Sacramento	6.5% (4;32)	9.5% (19;225)	24.3% (8;22)	5.2% (11;201)		1.7% (3;184)	3.3% (11;329)	3.5% (3;70)	2.7% (4;208)
	Sutter General Hospital	8.3% (9;68)	9.8% (38;362)	25.4% (22;74)	5.1% (12;266)	19.0% (4;22)	2.0% (5;234)	3.6% (8;233)	2.0% (4;196)	4.2% (8;218)
	Sutter Memorial Hospital	5.4% (24;487)	8.1% (10;111)	16.1% (5;25)	4.6% (3;84)		2.4% (2;76)	3.0% (13;430)		3.9% (4;96)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
STATEWIDE		6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Sacramento (continued)	UC Davis Medical Center	5.8% (23;402)	9.6% (43;410)	24.7% (20;75)	4.6% (14;291)	24.4% (9;44)	2.5% (7;297)	2.6% (13;602)	5.0% (3;70)	3.2% (11;388)
San Benito	Hazel Hawkins Memorial Hospital	4.2% (1;21)	11.1% (2;18)		4.9% (1;17)		2.9% (1;56)	2.0% (1;74)	0.0% (0;18)	17.2% (8;87) Worse
San Bernardino	Arrowhead Regional Medical Center	15.5% (12;107) Worse	11.5% (41;405)	29.8% (28;125)	3.0% (5;255)	39.8% (8;25)	5.7% (8;199) Worse	2.8% (7;584)	0.0% (0;42)	6.7% (6;230)
	Barstow Community Hospital	9.4% (3;33)	0.0% (0;60)		0.0% (0;60)		2.8% (1;38)	0.0% (0;129)		0.0% (0;70) Better
	Bear Valley Community Hospital							0.0% (0;8)		0.0% (0;29)
	Chino Valley Medical Center	3.6% (2;23)	2.9% (1;44)	0.0% (0;3)	2.4% (1;41)		0.9% (1;74)	3.2% (4;85)	0.0% (0;16)	0.0% (0;58)
	Colorado River Medical Center						0.0% (0;5)	0.0% (0;8)		0.0% (0;20)
	Community Hospital of San Bernardino	11.7% (4;32)	0.0% (0;17)		0.0% (0;17)		3.3% (3;84)	3.3% (2;116)	0.0% (0;19)	1.2% (1;128)
	Desert Valley Hospital	5.7% (15;204)	2.0% (2;148) Better	0.0% (0;5)	0.6% (1;142) Better		2.0% (4;162)	1.5% (7;439)	0.9% (1;65)	0.8% (2;253) Better
	Hi-Desert Medical Center	7.2% (1;11)	24.7% (3;32)		11.6% (2;31)		4.1% (1;41)	2.6% (2;109)	5.3% (1;24)	9.7% (12;164) Worse
	Kaiser Foundation Hospital – Fontana	6.2% (23;257)	9.5% (52;593)	19.2% (23;179)	5.8% (22;381)	28.5% (7;33)	2.2% (10;538)	2.5% (25;781)	2.8% (4;187)	3.4% (24;529)
	Loma Linda University Medical Center	5.6% (25;531)	9.2% (80;624)	19.9% (35;181)	5.5% (27;370)	23.4% (18;73)	1.9% (3;198)	7.3% (28;488) Worse	1.2% (1;104)	5.1% (11;224)
	Montclair Hospital Medical Center	2.9% (1;27)	35.2% (1;9)		21.2% (1;9)		5.0% (2;26)	0.0% (0;67)	3.0% (1;12)	0.0% (0;69)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
San Bernardino (continued)	Mountains Community Hospital						100.0% (1;3) Worse	0.0% (0;4)		20.9% (2;35)
	Redlands Community Hospital	5.2% (6;81)	11.0% (15;194)	13.1% (2;21)	6.8% (12;171)		1.9% (3;182)	2.6% (6;221)	1.3% (2;173)	4.1% (17;361)
	Saint Bernardine Medical Center	5.4% (34;996)	10.8% (15;178)	22.0% (7;46)	6.6% (6;126)	39.1% (2;6)	5.5% (10;186) Worse	2.5% (9;446)	0.0% (0;94)	4.1% (13;319)
	Saint Mary Regional Medical Center	6.1% (26;450)	7.2% (8;209)	0.0% (0;3)	4.2% (8;206)		1.7% (4;288)	1.9% (9;561)	1.1% (1;120)	4.6% (18;367)
	San Antonio Community Hospital	3.6% (17;487) Better	7.7% (26;358)	17.9% (13;77)	4.4% (11;275)	20.2% (2;6)	2.1% (7;289)	2.0% (10;459)	0.6% (1;152)	3.7% (21;522)
	Victor Valley Global Medical Center*	7.0% (2;35)	12.1% (4;63)		5.7% (2;61)		3.3% (2;64)	2.4% (3;203)	5.5% (2;43)	3.2% (4;139)
San Diego	Alvarado Hospital	9.8% (13;106)	12.0% (11;108)	16.7% (3;23)	8.1% (7;83)		2.4% (3;106)	1.1% (2;195)	2.1% (1;52)	1.4% (3;180) Better
	Fallbrook Hospital District	5.3% (1;13)	14.7% (2;23)	56.3% (1;4)	0.0% (0;18)		3.4% (1;43)	2.2% (1;70)	0.0% (0;36)	0.0% (0;69)
	Grossmont Hospital	5.9% (19;349)	10.4% (71;665)	25.4% (33;114)	5.5% (32;524)	25.0% (6;27)	3.0% (13;447)	3.6% (30;896)	3.2% (6;206)	3.0% (15;531)
	Kaiser Foundation Hospital – San Diego	8.5% (19;143)	10.9% (47;472)	28.6% (27;121)	4.8% (14;328)	31.4% (6;23)	3.3% (11;389)	4.2% (27;758)	2.3% (5;296)	6.3% (32;548) Worse
	Palomar Health Downtown Campus	9.3% (36;436) Worse	13.0% (88;630) Worse	32.3% (47;132) Worse	6.5% (27;461)	35.7% (14;37)	1.8% (5;351)	3.8% (26;665)	1.9% (3;182)	3.7% (23;578)
	Paradise Valley Hospital	3.4% (4;51)	3.5% (2;70)	10.3% (1;12)	1.6% (1;58)		3.0% (3;72)	1.8% (3;228)	4.8% (1;20)	4.3% (4;118)
	Pomerado Hospital	1.8% (1;33)	14.8% (10;110)	100.0% (2;4)	7.1% (8;105)		2.0% (3;147)	3.8% (10;187)	3.0% (3;84)	6.6% (17;166) Worse

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
STATEWIDE		6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
San Diego (continued)	Scripps Green Hospital	2.1% (3;204) Better	8.0% (22;193)	17.7% (9;52)	5.6% (8;107)	13.9% (5;34)	0.9% (1;146)	1.9% (6;286)	1.1% (1;93)	0.6% (1;133) Better
	Scripps Memorial Hospital – Encinitas (View comment letter)	2.9% (5;188)	8.6% (15;211)	25.1% (8;29)	4.1% (7;180)		2.8% (5;181)	2.0% (9;312)	1.4% (2;135)	3.7% (7;160)
	Scripps Memorial Hospital – La Jolla	5.6% (17;372)	11.5% (49;313)	23.0% (16;62)	7.3% (21;213)	28.6% (12;38)	1.4% (2;155)	4.1% (18;325)	2.8% (5;180)	4.8% (17;227)
	Scripps Mercy Hospital	6.1% (35;451)	10.0% (51;529)	24.3% (27;89)	5.7% (23;431)	15.2% (1;9)	1.0% (5;441) Better	2.3% (21;874)	2.2% (4;203)	2.6% (15;656)
	Sharp Chula Vista Medical Center	3.8% (6;162)	10.0% (28;314)	27.1% (11;38)	5.0% (12;260)	25.9% (5;16)	1.9% (4;230)	2.4% (19;714)	3.4% (5;115)	3.6% (13;363)
	Sharp Coronado Hospital and Healthcare Center	13.1% (1;3)	0.0% (0;19)		0.0% (0;19)		0.0% (0;33)	0.0% (0;50)	0.0% (0;20)	0.0% (0;17)
	Sharp Memorial Hospital	5.8% (15;306)	11.1% (43;475)	16.6% (13;89)	8.9% (29;366) Worse	13.8% (1;20)	2.2% (6;306)	5.0% (28;658) Worse	2.5% (7;288)	4.8% (22;535)
	Tri-City Medical Center – Oceanside	6.9% (21;302)	11.3% (45;386)	28.7% (26;84)	5.7% (16;289)	28.4% (3;13)	3.7% (12;273)	4.5% (16;382)	4.4% (7;188)	4.8% (12;320)
	UC San Diego Health System – Hillcrest Medical Center*	5.1% (12;286)	7.5% (36;341)	18.9% (22;94)	3.6% (10;227)	22.1% (4;20)	0.6% (2;348) Better	3.9% (19;656)	0.0% (0;100)	1.7% (4;316)
San Francisco	California Pacific Medical Center – Davies Campus	12.2% (1;3)	7.1% (29;341)	16.1% (9;54)	4.0% (14;237)	16.5% (6;50)	3.6% (1;33)	5.4% (1;31)	0.0% (0;10)	5.6% (3;59)
	California Pacific Medical Center – Pacific Campus	6.5% (15;244)	10.2% (35;288)	26.7% (22;62)	4.2% (10;223)	46.2% (3;3)	1.7% (7;316)	2.5% (11;390)	1.2% (2;156)	4.7% (13;288)
	California Pacific Medical Center – St. Luke’s Campus	0.0% (0;11)	12.8% (3;29)	34.5% (2;3)	6.0% (1;26)		1.4% (1;63)	3.1% (3;113)	0.0% (0;27)	2.3% (2;91)
	Chinese Hospital	0.0% (0;16)	3.7% (1;55)		2.1% (1;54)		0.0% (0;82)	1.7% (2;122)	4.0% (1;29)	5.8% (8;135)
	Kaiser Foundation Hospital – San Francisco	4.9% (24;628)	9.4% (21;220)	12.8% (4;28)	6.5% (16;189)	15.4% (1;3)	1.8% (3;182)	3.1% (13;320)	1.8% (1;58)	1.0% (1;83)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
San Francisco (continued)	Laguna Honda Hospital and Rehabilitation Center									0.0% (0;25)
	Saint Francis Memorial Hospital	21.0% (4;13) Worse	11.0% (12;110)	22.9% (5;21)	5.9% (6;87)		1.6% (2;101)	3.8% (5;142)	2.1% (1;53)	4.9% (7;201)
	Saint Mary's Medical Center, San Francisco	6.4% (5;110)	7.9% (8;105)	23.0% (4;16)	2.3% (2;86)	46.5% (2;3)	0.7% (1;129)	3.3% (9;214)	0.0% (0;55)	3.5% (7;158)
	San Francisco General Hospital	8.6% (13;175)	15.2% (46;264) Worse	31.1% (23;81)	9.4% (17;174) Worse	49.9% (6;9) Worse	1.8% (3;176)	2.3% (6;497)	1.7% (1;40)	2.5% (3;268)
	UC San Francisco Medical Center	11.0% (18;162) Worse	9.1% (72;481)	26.1% (35;125)	3.5% (10;194)	19.1% (27;162)	2.9% (7;209)	3.9% (14;373)	1.6% (1;60)	2.2% (5;272)
San Joaquin	Dameron Hospital	7.8% (13;277)	8.9% (5;94)	43.9% (2;4)	4.2% (3;89)		4.6% (6;148)	2.6% (5;215)	4.1% (2;47)	4.5% (10;213)
	Doctors Hospital of Manteca	7.9% (2;16)	9.4% (2;51)	42.0% (1;5)	3.5% (1;46)		0.0% (0;88)	2.7% (3;130)	4.3% (1;37)	11.2% (9;116) Worse
	Kaiser Foundation Hospital – Manteca	4.3% (4;81)	6.7% (5;98)	22.0% (2;8)	3.2% (3;89)		2.2% (4;142)	2.7% (11;284)	2.8% (4;134)	3.4% (3;64)
	Lodi Memorial Hospital	4.6% (3;50)	14.5% (6;70)	56.9% (3;7)	5.5% (3;63)		1.8% (3;143)	3.0% (7;284)	1.3% (1;94)	4.4% (9;288)
	Saint Joseph's Medical Center of Stockton	7.8% (26;370)	12.8% (46;313) Worse	24.5% (20;72)	9.4% (22;235) Worse	27.3% (4;6)	2.7% (7;297)	3.5% (17;530)	0.0% (0;130)	6.8% (24;420) Worse
	San Joaquin General Hospital	5.4% (2;45)	12.5% (16;147)	32.3% (13;35)	5.3% (3;111)		2.3% (2;136)	2.9% (4;209)	3.7% (1;27)	3.8% (2;87)
	Sutter Tracy Community Hospital	7.3% (2;24)	10.8% (3;41)		2.9% (1;38)		0.0% (0;79)	4.0% (4;103)	3.4% (1;27)	5.1% (4;86)
San Luis Obispo	French Hospital Medical Center	5.9% (13;295)	6.0% (2;52)		4.1% (2;51)		1.2% (1;76)	2.5% (4;129)	1.2% (1;50)	0.7% (1;100) Better

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
San Luis Obispo (continued)	Marian Regional Medical Center – Arroyo Grande*	3.9% (1;15)	18.8% (11;67) Worse	42.3% (6;10)	11.8% (5;57)		0.0% (0;65)	8.9% (8;88) Worse	9.7% (5;52) Worse	11.7% (8;71) Worse
	Sierra Vista Regional Medical Center	6.6% (2;13)	10.5% (13;128)	25.0% (8;49)	6.1% (5;78)		2.8% (1;67)	7.5% (5;53)	0.0% (0;52)	0.0% (0;82)
	Twin Cities Community Hospital	19.3% (5;17) Worse	16.9% (6;98)	90.1% (2;5)	7.3% (4;93)		1.7% (1;113)	2.2% (2;120)	5.8% (5;84) Worse	4.6% (8;220)
San Mateo	Kaiser Foundation Hospital – Redwood City	7.7% (5;71)	6.4% (53;461) Better	12.7% (27;190) Better	7.6% (16;191)	12.1% (10;80) Better	2.8% (2;92)	4.7% (10;174)	0.9% (1;62)	3.3% (4;105)
	Kaiser Foundation Hospital – South San Francisco	3.9% (3;39)	6.5% (11;154)	18.6% (6;31)	3.0% (5;120)	0.0% (0;3)	1.7% (3;140)	1.4% (5;245)	0.0% (0;97)	4.0% (5;106)
	Peninsula Medical Center	5.2% (12;239)	11.0% (31;271)	24.8% (14;49)	6.2% (16;218)	41.3% (1;4)	2.1% (4;215)	3.2% (15;360)	2.0% (4;178)	2.6% (7;276)
	San Mateo Medical Center	7.2% (1;17)	11.0% (5;33)	29.7% (5;10)	0.0% (0;22)		1.9% (1;102)	2.9% (2;119)	0.0% (0;13)	4.3% (2;110)
	Sequoia Hospital	10.9% (5;80)	9.7% (6;91)	22.7% (2;7)	5.6% (4;83)		3.0% (2;68)	1.4% (3;214)	0.9% (1;76)	6.4% (9;128)
	Seton Medical Center	4.6% (8;173)	8.8% (20;180)	21.3% (13;45)	5.6% (7;131)	0.0% (0;4)	1.5% (3;184)	3.4% (15;374)	1.7% (1;68)	2.0% (6;252)
Santa Barbara	Goleta Valley Cottage Hospital		30.9% (2;4)		16.3% (2;4)			17.5% (3;19) Worse	0.0% (0;61)	5.0% (1;36)
	Lompoc Valley Medical Center	20.6% (1;4)	15.7% (6;41)	32.7% (3;5)	10.6% (3;36)		6.0% (3;78)	11.4% (4;57) Worse	0.0% (0;30)	7.1% (7;135)
	Marian Regional Medical Center	8.2% (15;183)	14.1% (23;189) Worse	34.6% (12;41)	7.4% (11;148)		2.3% (5;226)	2.4% (7;248)	0.8% (1;110)	3.8% (8;150)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Santa Barbara (continued)	Santa Barbara Cottage Hospital	5.3% (9;255)	11.6% (68;567) Worse	20.4% (18;113)	8.5% (33;365) Worse	26.1% (17;89)	2.3% (6;279)	1.1% (4;357) Better	0.8% (1;157)	4.7% (15;362)
	Santa Ynez Valley Cottage Hospital		69.0% (1;5)		36.9% (1;5)			0.0% (0;17)		0.0% (0;23)
Santa Clara	El Camino Hospital	3.6% (7;219)	8.7% (27;276)	18.2% (12;60)	5.3% (11;208)	26.2% (4;8)	1.9% (5;301)	3.2% (16;390)	0.6% (1;207)	3.2% (11;344)
	Good Samaritan Hospital – San Jose	12.9% (22;242) Worse	11.2% (40;399)	27.5% (15;58)	6.0% (15;301)	25.8% (10;40)	4.3% (11;260)	6.0% (22;307) Worse	0.9% (1;147)	7.9% (19;274) Worse
	Kaiser Foundation Hospital – San Jose	7.3% (17;166)	7.5% (18;240)	13.2% (5;40)	5.2% (13;195)	0.0% (0;5)	0.7% (1;186)	2.3% (10;344)	5.3% (7;108) Worse	2.5% (4;131)
	Kaiser Foundation Hospital – Santa Clara	3.3% (19;973) Better	8.0% (25;337)	23.6% (15;57)	3.5% (9;274)	10.8% (1;6)	1.9% (6;271)	3.8% (34;733)	2.7% (7;190)	3.6% (9;196)
	O'Connor Hospital – San Jose	10.2% (20;213) Worse	11.7% (23;184)	23.4% (9;40)	7.7% (10;137)	28.4% (4;7)	4.3% (7;202)	3.7% (17;338)	3.7% (4;114)	4.4% (13;282)
	Regional Medical of San Jose	7.9% (33;301)	7.6% (51;476)	15.9% (24;137) Better	3.9% (14;301)	27.0% (13;38)	1.6% (6;312)	2.4% (12;522)	2.3% (2;86)	6.5% (23;316) Worse
	Saint Louise Regional Hospital	7.7% (3;18)	6.6% (2;56)	18.8% (1;8)	3.2% (1;48)		4.0% (2;42)	1.0% (1;111)	0.0% (0;38)	3.6% (4;112)
	Santa Clara Valley Medical Center	3.2% (5;227)	7.7% (19;352)	23.6% (15;75)	2.0% (3;268)	15.4% (1;9)	1.0% (2;321)	1.9% (6;643)	2.0% (1;43)	2.0% (3;257)
	Stanford Hospital	6.1% (13;228)	9.4% (53;465)	19.6% (18;103)	7.2% (25;268)	16.5% (10;94)	2.1% (6;332)	3.4% (17;412)	1.4% (2;131)	5.6% (18;379)
Santa Cruz	Dominican Hospital – Santa Cruz/Soquel	6.3% (11;242)	9.5% (18;214)	24.7% (9;44)	3.3% (4;162)	28.8% (5;8)	5.4% (7;182)	3.1% (8;329)	0.0% (0;118)	4.0% (6;220)
	Watsonville Community Hospital	11.2% (2;10)	11.2% (3;62)	34.8% (2;7)	4.2% (1;55)		3.1% (2;87)	4.8% (5;104)	2.6% (1;46)	3.8% (6;143)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
STATEWIDE		6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Shasta	Mayers Memorial Hospital		21.4% (1;5)		22.8% (1;4)		0.0% (0;3)	0.0% (0;21)		20.9% (3;27) Worse
	Mercy Medical Center – Redding	5.8% (12;255)	9.9% (27;266)	18.9% (12;70)	5.7% (11;187)	85.3% (4;9) Worse	4.3% (8;182)	3.5% (18;394)	1.8% (5;240)	4.0% (13;299)
	Shasta Regional Medical Center (View comment letter)	4.7% (7;182)	13.8% (9;136)	100.0% (3;5) Worse	5.4% (6;131)		0.7% (2;199)	2.8% (10;345)	1.4% (1;66)	5.2% (10;192)
Siskiyou	Fairchild Medical Center	13.4% (1;5)	9.7% (2;34)	34.9% (2;5)	0.0% (0;29)		0.0% (0;44)	5.2% (2;42)	5.4% (1;25)	4.6% (3;110)
	Mercy Medical Center Mt. Shasta	29.5% (4;7) Worse	25.4% (2;18)		8.4% (1;16)		0.0% (0;14)	11.3% (2;27)	6.5% (2;22)	24.7% (6;48) Worse
Solano	Kaiser Foundation Hospital – Rehabilitation Center Vallejo	4.4% (9;166)	5.8% (15;252) Better	17.0% (5;23)	2.0% (6;225) Better	44.2% (4;4)	1.6% (4;207)	2.3% (10;320)	1.5% (2;102)	2.6% (3;111)
	Kaiser Foundation Hospital – Vacaville	2.0% (2;72)	10.7% (11;116)	26.6% (3;11)	5.9% (8;104)		3.9% (4;107)	2.2% (4;138)	5.8% (5;74)	6.4% (6;96)
	North Bay Medical Center	4.1% (11;247)	12.6% (18;161)	33.4% (8;19)	6.3% (8;139)	33.1% (2;3)	1.8% (4;173)	3.6% (11;328)	1.3% (1;63)	3.0% (6;220)
	Sutter Solano Medical Center	5.3% (2;13)	0.0% (0;87) Better		0.0% (0;86) Better		3.1% (3;86)	3.4% (4;151)	0.0% (0;32)	2.9% (3;110)
Sonoma	Healdsburg District Hospital		0.0% (0;23)		0.0% (0;22)		0.0% (0;7)	0.0% (0;21)	0.0% (0;17)	1.9% (1;48)
	Kaiser Foundation Hospital – Santa Rosa	0.9% (1;85) Better	10.3% (22;173)	22.0% (8;28)	6.0% (13;143)		2.3% (3;173)	1.5% (3;175)	1.2% (1;108)	2.9% (3;111)
	Palm Drive Hospital		0.0% (0;34)	0.0% (0;3)	0.0% (0;31)		0.0% (0;16)	3.2% (1;34)	0.0% (0;17)	7.5% (4;51)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see [Technical Note](#) for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Sonoma (continued)	Petaluma Valley Hospital	4.6% (2;32)	10.1% (4;38)	28.3% (3;7)	4.0% (1;31)		3.5% (2;56)	4.3% (4;100)	0.0% (0;34)	1.2% (1;87)
	Santa Rosa Memorial Hospital – Montgomery	7.6% (15;266)	11.5% (28;271)	26.6% (15;70)	6.7% (13;199)		2.1% (5;234)	6.3% (22;281) Worse	2.4% (4;157)	7.1% (17;212) Worse
	Sonoma Valley Hospital	0.0% (0;10)	7.1% (3;39)	24.8% (3;6)	0.0% (0;33)		0.0% (0;27)	4.5% (2;30)	2.4% (1;34)	0.0% (0;21)
	Sutter Medical Center of Santa Rosa	5.1% (7;163)	18.4% (6;71)	68.8% (4;8) Worse	5.7% (2;63)		1.3% (1;89)	3.1% (5;154)	2.7% (1;41)	2.1% (2;86)
Stanislaus	Central Valley Specialty Hospital		24.3% (1;4)		13.1% (1;4)					31.2% (1;6)
	Doctors Medical Center	11.3% (29;375) Worse	12.1% (47;390) Worse	29.1% (29;95)	6.2% (15;290)	41.3% (3;5)	6.1% (13;272) Worse	2.6% (9;498)	0.9% (1;136)	5.2% (19;472)
	Emanuel Medical Center, Inc.	5.4% (14;295)	9.5% (3;66)	21.1% (1;5)	5.4% (2;61)		1.6% (2;153)	5.4% (16;330) Worse	3.7% (1;35)	4.3% (9;272)
	Memorial Hospital Medical Center – Modesto	6.5% (27;426)	6.1% (20;358) Better	15.0% (12;83)	3.5% (8;269)	0.0% (0;6)	2.0% (8;357)	2.8% (13;504)	2.9% (7;221)	5.2% (23;485)
	Oak Valley District Hospital	14.6% (1;7)	0.0% (0;17)		0.0% (0;15)		0.0% (0;27)	2.5% (1;38)	8.6% (1;16)	5.0% (2;46)
Tehama	Saint Elizabeth Community Hospital	0.0% (0;4)	6.8% (3;45)		1.8% (1;42)		2.9% (2;72)	1.1% (1;89)	8.6% (1;12)	6.0% (7;142)
Trinity	Trinity Hospital	0.0% (0;3)					0.0% (0;6)	6.3% (1;22)		0.0% (0;36)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Tulare	Kaweah Delta Medical Center (View comment letter)	5.1% (20;443)	10.5% (22;318)	24.0% (9;31)	5.9% (12;283)	41.7% (1;4)	2.0% (7;365)	3.9% (18;542)	4.1% (6;161)	3.1% (21;678)
	Sierra View District Hospital	15.0% (10;59) Worse	12.7% (8;102)	32.4% (2;8)	6.8% (6;94)		4.8% (5;126)	12.0% (17;213) Worse	0.0% (0;66)	8.3% (21;275) Worse
	Tulare Regional Medical Center	0.0% (0;32)	9.6% (5;38)	27.9% (4;6)	3.3% (1;32)		4.5% (3;62)	4.5% (4;105)	5.4% (2;32)	6.2% (8;177)
Tuolumne	Sonora Regional Medical Center – Greenley	6.9% (3;22)	10.6% (8;89)	26.6% (3;8)	5.3% (4;80)		2.5% (2;86)	3.3% (5;125)	4.8% (4;66)	0.6% (1;176) Better
Ventura	Community Memorial Hospital – San Buenaventura	6.3% (15;241)	7.4% (12;225)	18.2% (5;33)	3.3% (5;186)	29.2% (2;6)	1.5% (3;186)	1.8% (6;328)	4.9% (5;122)	4.4% (13;249)
	Los Robles Hospital and Medical Center	6.8% (18;352)	10.8% (21;206)	21.7% (9;53)	6.2% (9;150)	43.8% (3;3)	1.2% (3;248)	2.2% (8;310)	1.2% (2;162)	2.9% (8;242)
	Ojai Valley Community Hospital	0.0% (0;7)	12.0% (1;15)	72.0% (1;3)	0.0% (0;11)		0.0% (0;16)	0.0% (0;25)	0.0% (0;13)	2.9% (2;46)
	Saint John’s Pleasant Valley Hospital	11.1% (3;17)	15.6% (7;79)	46.8% (3;8)	7.4% (4;71)		1.0% (1;74)	5.8% (12;137) Worse	4.5% (3;66)	5.8% (8;139)
	Saint John’s Regional Medical Center	4.7% (15;251)	11.4% (33;227)	30.7% (19;59)	5.0% (8;155)	27.4% (6;13)	2.8% (7;144)	3.9% (17;338)	3.9% (2;57)	4.2% (18;279)
	Simi Valley Hospital and Healthcare Services – Sycamore	5.8% (2;23)	17.4% (16;97) Worse	47.2% (9;21) Worse	8.1% (6;75)		5.1% (6;103)	5.4% (6;99)	0.0% (0;54)	7.0% (10;154)
	Ventura County Medical Center	5.9% (2;44)	14.0% (18;129)	33.8% (10;35)	7.0% (7;91)	53.8% (1;3)	3.2% (3;102)	1.1% (1;162)	2.9% (1;42)	5.0% (5;136)
Yolo	Sutter Davis Hospital	13.3% (1;4)	6.7% (2;41)		2.0% (1;39)		0.0% (0;47)	1.5% (1;46)	2.5% (1;40)	6.3% (3;59)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2013

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.										
	STATEWIDE	6.0% (2,803;46,984)	9.2% (5,072;54,983)	21.9% (2,355;10,730)	4.9% (2,046;41,526)	24.6% (671;2,727)	2.3% (1,087;47,723)	3% (2,510;83,220)	2.0% (494;24,103)	3.9% (2,487;63,853)
Yolo (continued)	Woodland Memorial Hospital	6.4% (3;22)	22.2% (10;56) Worse	27.5% (2;6)	17.1% (6;48) Worse		1.8% (1;55)	0.0% (0;45)	3.1% (1;38)	5.8% (4;89)
Yuba	Rideout Memorial Hospital	6.6% (16;263)	9.9% (23;234)	28.9% (12;37)	5.1% (10;190)	10.5% (1;7)	1.7% (4;274)	3.5% (14;422)	2.6% (3;120)	5.0% (16;377)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.5, and OSHPD hospital patient discharge data, 2013. See [Technical Note](#) for details about methodology.

*Hospital name changed from 2012 to 2013, see Technical Note for details.

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.