



**PQI 01 and PQI 14: Diabetes Short-Term Complications and Uncontrolled
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 02: Perforated Appendix [ruptured appendix]
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 03: Diabetes Long-Term Complications
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 04: Pediatric Asthma
North Central Counties**




	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
North Central Counties	58.5	75.8	41.2	65.1	61.7	48	41.3	51.4	37.5	23.8
Statewide	126.9	131	109.6	117.7	113.3	98.2	90.1	87.3	78.5	77.6

**PQI 05: Chronic Obstructive Pulmonary Disease (COPD) [*chronic bronchitis or emphysema*]
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 06: Pediatric Gastroenteritis [severe vomiting and/or severe diarrhea in children]
North Central Counties**




	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
North Central Counties	93.1	97.4	100.3	77.2	116.9	100.7	132.3	91.8	73.4	89.8
Statewide	137.9	149.6	137.7	140	129.5	125.2	120	114.4	99.5	75.3

**PQI 07: Hypertension [*high blood pressure*]
North Central Counties**




**PQI 08: Congestive Heart Failure (CHF) [*heart failure*]
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 10: Dehydration
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 11: Bacterial Pneumonia
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 12: Urinary Tract Infection (UTI)
North Central Counties**




Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

**PQI 13: Angina without Procedure [chest pain]
North Central Counties**




	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
North Central Counties	43.8	74.4	59.1	39.3	31.3	24.3	23.3	21.3	9.1	11.5
Statewide	65.7	60.6	54.3	48.5	41.8	37.8	33	30.7	28	26

**PQI 15: Adult Asthma
North Central Counties**



Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2

PQI 16: Lower-Extremity Amputation among Patients with Diabetes
[removal of leg or foot due to diabetes complications]
North Central Counties



Source: Office of Statewide Health Planning and Development, Patient Discharge Data, 1999-2008
 Agency for Healthcare Research and Quality, Prevention Quality Indicators, Version 3.2