

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
Alameda	Alameda County Medical Center – Highland Campus				11.5% (3; 52)		0% (0; 3)	0% (0; 28)	1.7% (3; 357)	11.3% (20; 148)	2.5% (2; 120)	5.5% (1; 17)	5.8% (8; 213)
	Alameda Hospital							4.9% (5; 62)	1.7% (3; 150)	9.1% (3; 67)	2.3% (2; 91)	0% (0; 39)	5.2% (7; 122)
	Alta Bates Summit Medical Center – Alta Bates Campus				8.2% (8; 73)	3.4% (3; 91)	0% (0; 20)	7.1% (9; 122)	2.7% (13; 364)	9.8% (41; 312)	1.9% (3; 195)	2.6% (3; 85)	4.5% (11; 277)
	Alta Bates Summit Medical Center – Summit Campus – Hawthorne		9.3% (1; 5)	0% (0; 5)		2.1% (12; 1,117)	2.6% (1; 23)	6.5% (30; 820)	2.6% (15; 595)	10.5% (21; 253)	2.4% (7; 299)	2.4% (3; 99)	2.8% (13; 413) Better
	Eden Medical Center				7.9% (7; 62)		0% (0; 4)	5.3% (5; 53)	1.9% (6; 213)	8.4% (30; 185)	2.3% (4; 127)	2.1% (2; 81)	0.8% (2; 211) Better
	Kaiser Foundation Hospital – Fremont				0% (0; 4)			4.1% (6; 101)	2.6% (7; 212)	7.9% (11; 132)	1.9% (2; 146)	5.3% (2; 48)	3.3% (12; 223)
	Kaiser Foundation Hospital – Hayward	0% (0; 6)	13.4% (2; 13)	0% (0; 25)			2.8% (1; 45)	5.8% (16; 146)	3.7% (11; 258)	10.5% (22; 170)	1.4% (3; 235)	7.0% (4; 83)	4.3% (19; 330)
	Kaiser Foundation Hospital – Oakland Campus			0% (0; 20)			0% (0; 33)	7.1% (13; 135)	3.4% (18; 507)	9.3% (15; 229)	1.0% (2; 236)	2.3% (2; 134)	7.9% (27; 323) Worse
	Saint Rose Hospital					1.5% (2; 91)	0% (0; 3)	8.6% (10; 130)	3.3% (6; 204)	6.4% (4; 76)	0.8% (1; 166)	4.1% (1; 40)	1.8% (2; 171)
	San Leandro Hospital			0% (0; 13)			0% (0; 32)	3.1% (2; 31)	2.7% (8; 258)	9.2% (8; 96)	3.0% (5; 116)	0% (0; 33)	3.5% (6; 178)
	Valleycare Medical Center			11.4% (1; 9)		3.4% (6; 159)	0% (0; 30)	9.3% (16; 183)	5.8% (8; 156)	7.2% (6; 85)	1.2% (1; 114)	5.6% (2; 51)	6.0% (12; 236)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Washington Hospital – Fremont				9.0% (7; 53)	1.8% (10; 427)	0% (0; 37)	8.5% (18; 256)	3.4% (13; 379)	10.0% (38; 300)	4.3% (9; 257)	5.1% (4; 93)	7.6% (22; 305) Worse
Amador	Sutter Amador Hospital							0% (0; 11)	5.2% (5; 84)	14.2% (4; 54)	1.4% (1; 61)	0% (0; 40)	7.9% (6; 109)
Butte	Biggs Gridley Memorial Hospital								6.5% (2; 33)	32.8% (2; 8)	0% (0; 5)	19.2% (1; 12)	14.1% (6; 81) Worse
	Enloe Medical Center – Esplanade Campus			0% (0; 26)	3.0% (1; 39)	4.8% (12; 365) Worse	0% (0; 112)	12.9% (25; 281) Worse	2.2% (6; 272)	8.3% (17; 237)	3.0% (5; 216)	3.7% (6; 167)	7.4% (27; 562) Worse
	Feather River Hospital						0% (0; 18)	2.1% (2; 55) Better	0.8% (1; 94)	6.7% (6; 75)	0% (0; 104)	6.5% (4; 58)	2.4% (3; 106)
	Oroville Hospital				135.5% (1; 4)		0% (0; 21)	8.1% (8; 54)	2.8% (6; 238)	9.9% (9; 98)	1.9% (3; 165)	1.0% (1; 56)	3.4% (9; 313)
Calaveras	Mark Twain Saint Joseph’s Hospital							8.4% (1; 8)	2.8% (2; 69)	3.8% (1; 28)	0% (0; 38)	7.8% (2; 24)	10.4% (9; 87) Worse
Colusa	Colusa Regional Medical Center							19.1% (2; 6)	0% (0; 44)	0% (0; 10)	5.6% (1; 25)		2.8% (2; 62)
Contra Costa	Contra Costa Regional Medical Center				0% (0; 5)			18.5% (2; 16)	2.3% (4; 256)	3.5% (1; 55)	3.3% (2; 63)	5.6% (1; 20)	2.5% (3; 210)
	Doctors Medical Center – San Pablo			0% (0; 3)		2.5% (4; 205)	0% (0; 9)	4.0% (8; 189) Better	2.5% (9; 392)	5.6% (5; 146)	1.0% (1; 142)	3.7% (3; 67)	4.3% (11; 248)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	John Muir Medical Center – Concord Campus			0% (0; 30)		1.8% (9; 423)	0% (0; 56)	9.4% (29; 346)	3.3% (18; 363)	10.9% (18; 150)	2.2% (6; 220)	0% (0; 60)	5.0% (15; 240)
	John Muir Medical Center – Walnut Creek Campus (View Comment Letter)			10.9% (1; 5)	7.5% (12; 106)	1.0% (2; 199)	0% (0; 54)	5.1% (16; 211)	1.3% (8; 334) Better	9.7% (81; 363)	1.5% (4; 244)	1.7% (6; 209)	3.0% (15; 319) Better
	Kaiser Foundation Hospital – Antioch		0% (0; 4)					7.7% (7; 77)	4.0% (13; 233)	10.2% (13; 126)	3.6% (3; 109)	0% (0; 62)	5.9% (12; 193)
	Kaiser Foundation Hospital – Richmond Campus							6.2% (6; 83)	2.9% (6; 142)	12.0% (15; 101)	7.9% (4; 79)		6.2% (6; 88)
	Kaiser Foundation Hospital – Walnut Creek	13.7% (1; 7)	0% (0; 18)	7.9% (1; 28)		3.4% (2; 57)	0% (0; 63)	9.6% (17; 156)	4.9% (23; 340)	14.3% (34; 238) Worse	3.6% (13; 258)	4.6% (13; 207) Worse	5.4% (27; 409)
	San Ramon Regional Medical Center	0% (0; 9)				2.9% (1; 64)	0% (0; 7)	11.2% (5; 55)	2.1% (2; 80)	15.2% (8; 47)	2.6% (2; 79)	5.1% (2; 46)	6.7% (10; 177)
	Sutter Delta Medical Center							8.2% (10; 75)	1.7% (5; 236)	6.9% (6; 109)	2.3% (5; 150)	5.0% (3; 56)	4.7% (10; 207)
Del Norte	Sutter Coast Hospital							6.4% (5; 33)	0% (0; 72) Better	20.7% (8; 56)	0% (0; 71)		2.1% (3; 90)
El Dorado	Barton Memorial Hospital						0% (0; 6)	14.1% (2; 5)	2.8% (1; 43)	20.7% (3; 28)	2.4% (1; 55)	0% (0; 23)	4.2% (3; 91)
	Marshall Medical Center							9.1% (12; 55)	3.1% (7; 179)	15.6% (15; 96)	2.4% (3; 142)	8.1% (5; 70) Worse	3.6% (8; 208)
Fresno	Clovis Community Medical Center**					2.2% (1; 20)		13.5% (12; 91)	4.5% (8; 151)	13.7% (4; 44)	3.3% (3; 129)	3.3% (2; 62)	8.0% (13; 216)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Coalinga Regional Medical Center								9.0% (3; 41)	17.6% (1; 13)	0% (0; 3)		3.0% (1; 156)
	Community Regional Medical Center – Fresno		0% (0; 4)	0% (0; 9)	11.3% (16; 161)	3.2% (13; 600)	0% (0; 11)	8.4% (47; 640)	4.1% (30; 843)	9.4% (72; 553)	2.6% (15; 491)	3.6% (4; 120)	4.3% (26; 744)
	Fresno Heart and Surgical Hospital	20.3% (1; 3)	0% (0; 3)	13.4% (1; 13)		1.2% (2; 547)	0% (0; 167)	6.8% (7; 137)	3.7% (4; 143)	83.2% (1; 5)	0% (0; 8)		0% (0; 5)
	Kaiser Foundation Hospital – Fresno			0% (0; 8)			0% (0; 50)	5.9% (15; 163)	3.7% (17; 315)	8.3% (16; 151)	1.2% (1; 93)	3.3% (4; 107)	4.2% (17; 290)
	Kingsburg Medical Center												30.6% (1; 3)
	Saint Agnes Medical Center	9.5% (1; 6)	0% (0; 4)	5.4% (1; 22)	8.1% (5; 95)	1.7% (12; 633)	3.1% (1; 53)	7.3% (40; 539)	4.6% (42; 904)	12.6% (45; 442)	4.7% (23; 500) Worse	1.9% (4; 293)	8.3% (51; 797) Worse
	Sierra Kings District Hospital							0% (0; 7)	8.1% (2; 35)	18.9% (1; 13)	0% (0; 13)		7.4% (6; 126)
Glenn	Glenn Medical Center								11.3% (1; 10)	0% (0; 3)	0% (0; 6)		0% (0; 27)
Humboldt	Jerold Phelps Community Hospital								0% (0; 6)				0% (0; 19)
	Mad River Community Hospital						0% (0; 17)	38.8% (5; 8) Worse	9.7% (4; 53)	19.1% (5; 39)	4.1% (2; 49)	4.5% (1; 19)	4.2% (4; 113)
	Redwood Memorial Hospital							23.5% (5; 9) Worse	11.3% (3; 24)	11.1% (2; 19)	2.6% (1; 35)	0% (0; 36)	2.7% (2; 91)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Saint Joseph Hospital – Eureka			6.2% (1; 20)	0% (0; 16)	5.4% (5; 234) Worse	0% (0; 91)	13.8% (14; 140) Worse	3.7% (6; 98)	8.9% (12; 106)	4.6% (5; 85)	0% (0; 68)	4.9% (6; 117)
Imperial	El Centro Regional Medical Center				0% (0; 6)			13.0% (7; 40)	4.5% (9; 181)	17.0% (15; 92) Worse	2.7% (5; 149)	4.1% (2; 73)	3.8% (10; 169)
	Pioneers Memorial Hospital				0% (0; 8)			8.1% (4; 32)	4.8% (9; 152)	6.3% (3; 50)	6.8% (7; 107) Worse	5.1% (1; 25)	4.7% (10; 178)
Inyo	Northern Inyo Hospital							18.4% (1; 5)	10.1% (2; 21)	80.3% (4; 13) Worse	4.5% (1; 27)	0% (0; 19)	4.9% (2; 58)
Kern	Bakersfield Heart Hospital			0% (0; 15)		1.1% (7; 682)	0% (0; 26)	3.1% (10; 466) Better	0.9% (3; 271) Better	2.7% (1; 46)	0% (0; 66)		3.3% (6; 173)
	Bakersfield Memorial Hospital		0% (0; 3)	0% (0; 26)	9.2% (11; 92)	0.5% (2; 530) Better	0% (0; 41)	5.6% (15; 330)	1.8% (7; 388)	7.2% (17; 227) Better	1.1% (2; 205)	0% (0; 135)	4.0% (14; 408)
	Delano Regional Medical Center (View Comment Letter)							19.0% (5; 15)	5.4% (7; 124)	16.9% (5; 31)	8.0% (4; 60) Worse	15.6% (1; 12)	5.5% (14; 196)
	Good Samaritan Hospital – Bakersfield								4.4% (1; 36)		0% (0; 13)		2.6% (1; 94)
	Kern Medical Center				17.3% (7; 58) Worse			6.5% (1; 12)	0% (0; 92)	15.4% (11; 71)	0% (0; 89)	3.2% (1; 26)	4.3% (3; 131)
	Kern Valley Healthcare District							12.5% (3; 16)	2.5% (2; 65)	0% (0; 11)	0% (0; 13)		2.5% (2; 134)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Mercy Hospital – Bakersfield			0% (0; 11)	6.6% (8; 85)		0% (0; 35)	4.4% (4; 41)	0.8% (2; 208) Better	8.0% (25; 267)	2.0% (5; 258)	0.6% (1; 145) Better	4.5% (24; 593)
	Ridgecrest Regional Hospital							14.6% (1; 7)	8.6% (8; 123) Worse	11.6% (4; 44)	9.2% (3; 48)	0% (0; 3)	14.2% (9; 106) Worse
	San Joaquin Community Hospital		32.5% (1; 3)		10.9% (14; 82)	3.1% (5; 187)	0% (0; 8)	9.7% (11; 143)	2.6% (8; 360)	12.5% (45; 311)	2.9% (5; 195)	2.4% (2; 99)	7.5% (27; 362) Worse
	Tehachapi Hospital								0% (0; 13)				0% (0; 22)
Kings	Central Valley General Hospital								0% (0; 54)	5.1% (2; 24)	0% (0; 31)		4.5% (3; 82)
	Corcoran District Hospital								0% (0; 11)				0% (0; 15)
	Hanford Community Medical Center						0% (0; 5)	12.3% (12; 70)	6.7% (14; 210) Worse	13.2% (11; 86)	4.0% (7; 187)	1.8% (1; 80)	7.1% (23; 377) Worse
Lake	Redbud Community Hospital**							20.6% (2; 4)	4.3% (1; 38)	22.4% (1; 10)	6.2% (3; 59)	0% (0; 14)	7.9% (6; 137)
	Sutter Lakeside Hospital							5.5% (1; 9)	0% (0; 48)	4.7% (2; 49)	3.3% (3; 53)	0% (0; 34)	4.2% (3; 76)
Lassen	Banner Lassen Medical Center								10.7% (1; 14)	25.3% (1; 11)	8.8% (1; 16)		9.2% (3; 67)
Los Angeles	Alhambra Hospital				0% (0; 3)			7.3% (8; 44)	1.4% (4; 203)	6.3% (5; 74)	1.7% (3; 115)	3.4% (1; 38)	4.0% (13; 219)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Antelope Valley Hospital		76.5% (1; 3)	0% (0; 3)	9.9% (11; 50)	3.1% (10; 288)	0% (0; 39)	8.2% (25; 349)	3.1% (16; 536)	8.1% (32; 311) Better	3.8% (13; 319)	2.4% (4; 153)	4.7% (31; 595)
	Bellflower Medical Center							7.2% (1; 13)	1.0% (1; 98)	0% (0; 24)	2.9% (3; 63)	0% (0; 10)	2.4% (2; 110)
	Beverly Hospital				5.2% (1; 9)	2.9% (9; 129)	2.6% (1; 25)	9.5% (19; 142)	1.8% (8; 388) Better	9.4% (20; 176)	1.4% (3; 179)	0.9% (1; 72)	6.2% (20; 240)
	Brotman Medical Center			0% (0; 12)	4.3% (1; 70)	1.6% (1; 77)	0% (0; 13)	8.5% (9; 88)	5.0% (13; 223)	8.3% (11; 141)	2.3% (4; 133)	2.7% (2; 35)	4.8% (15; 249)
	California Hospital Medical Center – Los Angeles				14.4% (6; 19)		0% (0; 5)	8.9% (12; 95)	2.2% (7; 469)	10.2% (28; 235)	1.9% (3; 203)	2.2% (1; 43)	3.9% (6; 182)
	Catalina Island Medical Center												0% (0; 3)
	Cedars Sinai Medical Center	6.3% (1; 14)	0% (0; 28)	0% (0; 40)	7.9% (34; 542)	1.7% (15; 919)	0% (0; 122)	8.8% (43; 510)	3.2% (45; 1,141)	8.0% (55; 530) Better	2.5% (13; 478)	1.6% (5; 221)	4.6% (38; 837)
	Centinela Hospital Medical Center (View Comment Letter)			0% (0; 6)	11.8% (36; 80) Worse	2.2% (6; 269)	0% (0; 20)	5.8% (24; 327)	0.9% (11; 1,140) Better	6.8% (64; 555) Better	0.2% (1; 370) Better	0% (0; 73)	2.6% (17; 360) Better
	Citrus Valley Medical Center – Inter Community Campus			0% (0; 4)	0% (0; 14)	2.4% (6; 272)	0% (0; 50)	8.9% (24; 330)	3.6% (14; 358)	7.8% (12; 145)	2.2% (3; 137)	4.8% (2; 58)	3.3% (8; 201)
	Citrus Valley Medical Center – QV Campus			22.5% (2; 9) Worse	6.3% (2; 32)		0% (0; 15)	5.4% (6; 67)	2.2% (7; 312)	7.7% (17; 193)	1.4% (2; 165)	4.4% (3; 64)	5.1% (8; 171)
	City of Hope Helford Clinical Research Hospital	0% (0; 13)	7.2% (1; 18)		2.8% (2; 102)				62.9% (2; 8) Worse	10.2% (1; 10)	0% (0; 13)		12.0% (9; 106) Worse

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Coast Plaza Doctors Hospital							7.1% (1; 13)	0.6% (1; 139) Better	7.4% (2; 22)	4.6% (2; 48)	0% (0; 18)	4.4% (8; 151)
	Community Hospital of Long Beach							8.8% (10; 48)	0% (0; 98)	9.0% (4; 42)	2.4% (1; 56)	0% (0; 29)	4.2% (9; 145)
	Community and Mission Hospital of Huntington Park – Slauson							16.3% (1; 5)	0.9% (1; 149)	0% (0; 21)	0% (0; 58)	0% (0; 6)	6.3% (5; 176)
	Doctors Hospital of West Covina, Inc.										0% (0; 4)		0% (0; 3)
	Downey Regional Medical Center			0% (0; 5)	7.3% (3; 70)	2.6% (8; 295)	0% (0; 13)	7.2% (22; 216)	2.9% (17; 558)	8.0% (22; 237)	0.7% (2; 265)	0% (0; 119)	2.6% (11; 381) Better
	East Los Angeles Doctors Hospital							5.4% (4; 39)	2.8% (4; 92)	4.3% (1; 45)	4.0% (2; 48)	0% (0; 6)	2.7% (6; 105)
	East Valley Hospital Medical Center							2.7% (1; 17)	0% (0; 20)	14.0% (1; 22)	0% (0; 18)	0% (0; 6)	0.7% (1; 92) Better
	Encino Hospital Medical Center							6.4% (4; 26)	1.5% (3; 90)	8.6% (4; 42)	1.5% (2; 66)	4.0% (4; 86)	3.0% (6; 160)
	Foothill Presbyterian Hospital – Johnston Memorial				8.0% (1; 11)		0% (0; 8)	10.0% (7; 44)	2.4% (9; 283)	8.1% (9; 112)	3.2% (4; 126)	0% (0; 74)	3.6% (8; 172)
	Garfield Medical Center				10.0% (6; 33)	1.2% (6; 223)	0% (0; 20)	5.8% (13; 174)	2.2% (9; 352)	7.4% (22; 255) Better	2.2% (5; 197)	0% (0; 77)	6.0% (29; 345)
	Glendale Adventist Medical Center – Wilson Terrace			0% (0; 9)	7.3% (17; 145)	2.0% (12; 644)	0% (0; 16)	6.8% (19; 296)	3.1% (24; 441)	8.0% (41; 328) Better	2.6% (7; 230)	5.5% (8; 119)	5.8% (36; 394)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Glendale Memorial Hospital and Medical Center		15.7% (1; 4)	0% (0; 11)	11.6% (4; 14)	2.4% (11; 388)	0% (0; 25)	7.7% (27; 324)	1.9% (13; 539) Better	7.9% (14; 169)	1.7% (4; 206)	2.6% (4; 95)	3.4% (10; 301)
	Good Samaritan Hospital – Los Angeles		0% (0; 6)	0% (0; 13)	19.6% (4; 38)	2.9% (19; 547) Worse	0% (0; 13)	8.2% (24; 325)	3.8% (23; 645)	9.0% (22; 208)	2.4% (5; 218)	3.9% (3; 77)	3.9% (15; 299)
	Greater El Monte Community Hospital							4.2% (2; 21)	0% (0; 83)	0% (0; 51) Better	4.5% (3; 41)	0% (0; 7)	0% (0; 82) Better
	Harbor – UCLA Medical Center	0% (0; 3)	0% (0; 6)	0% (0; 13)	10.2% (5; 101)	4.2% (10; 396) Worse	124.5% (1; 5) Worse	8.4% (10; 257)	1.9% (5; 489)	9.3% (18; 239)	2.4% (4; 170)	2.8% (1; 25)	6.0% (13; 300)
	Henry Mayo Newhall Memorial Hospital				9.8% (4; 40)		0% (0; 22)	5.5% (12; 118)	3.6% (14; 231)	9.2% (16; 178)	2.2% (4; 165)	3.7% (4; 95)	3.0% (9; 244)
	Hollywood Community Hospital							12.1% (1; 10)	1.4% (1; 79)	6.2% (1; 27)	2.7% (2; 46)		0% (0; 94)
	Hollywood Presbyterian Medical Center				5.0% (1; 10)		0% (0; 5)	8.9% (17; 117)	4.3% (21; 447)	9.9% (21; 167)	2.6% (6; 212)	5.7% (4; 60)	4.8% (17; 266)
	Huntington Memorial Hospital	0% (0; 3)	0% (0; 8)	7.2% (1; 15)	7.4% (8; 183)	1.3% (5; 395)	5.9% (1; 35)	6.3% (18; 269)	1.9% (12; 566) Better	8.7% (35; 376)	1.9% (5; 317)	0.6% (1; 187)	2.8% (13; 436) Better
	Kaiser Foundation Hospital – Baldwin Park			0% (0; 15)			0% (0; 58)	9.2% (11; 95)	2.1% (12; 492)	8.2% (20; 261)	1.7% (5; 311)	2.6% (2; 78)	2.0% (10; 432) Better
	Kaiser Foundation Hospital – Bellflower			0% (0; 27)			0% (0; 48)	6.5% (14; 156)	2.2% (13; 537)	8.2% (19; 269)	2.1% (7; 312)	1.1% (1; 93)	2.8% (11; 425)
	Kaiser Foundation Hospital – Harbor City	0% (0; 4)	23.2% (1; 7)	0% (0; 3)		0% (0; 3)	0% (0; 50)	2.8% (4; 94) Better	2.5% (15; 546)	8.1% (12; 212)	1.9% (6; 347)	1.8% (1; 54)	3.5% (22; 523)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Kaiser Foundation Hospital – Panorama City	0% (0; 3)	0% (0; 6)	30.0% (1; 4)		0% (0; 10)	0% (0; 34)	7.2% (9; 99)	2.2% (13; 462)	8.4% (12; 210)	1.8% (5; 263)	3.2% (3; 130)	3.3% (16; 401)
	Kaiser Foundation Hospital – Sunset	0% (0; 3)	0% (0; 9)	0% (0; 38)	6.8% (22; 376)	1.9% (8; 541)	2.5% (1; 25)	7.6% (51; 950)	2.1% (16; 522) Better	10.3% (66; 491)	1.7% (6; 263)	0% (0; 67)	2.6% (12; 350) Better
	Kaiser Foundation Hospital – West Los Angeles	0% (0; 7)					0% (0; 12)	8.0% (17; 129)	1.3% (6; 397) Better	8.9% (21; 270)	2.5% (8; 307)	0% (0; 44)	5.0% (29; 364)
	Kaiser Foundation Hospital – Woodland Hills			16.8% (2; 10) Worse	3.8% (3; 57)		0% (0; 83)	6.0% (13; 116)	3.2% (24; 385)	6.9% (27; 281) Better	1.5% (6; 275)	2.7% (6; 152)	2.1% (10; 346) Better
	Lakewood Regional Medical Center				12.5% (7; 28)	1.7% (3; 297)	0% (0; 24)	7.4% (13; 196)	2.6% (12; 501)	8.9% (25; 264)	2.7% (5; 179)	6.7% (4; 94)	5.7% (14; 247)
	Lancaster Community Hospital			0% (0; 6)	9.9% (3; 19)	0% (0; 189)	6.8% (1; 18)	2.7% (3; 151) Better	2.2% (7; 354)	8.0% (9; 114)	2.9% (4; 140)	2.7% (1; 63)	5.2% (11; 246)
	Little Company of Mary Medical Center – San Pedro**			0% (0; 4)	0% (0; 22)		0% (0; 9)	12.3% (4; 17)	2.0% (4; 156)	11.6% (12; 96)	2.6% (3; 99)	0% (0; 61)	4.5% (11; 194)
	Little Company of Mary Medical Center – Torrance**			0% (0; 6)	11.3% (3; 27)	2.7% (12; 495)	0% (0; 46)	8.4% (21; 235)	3.4% (25; 635)	10.2% (27; 263)	1.0% (3; 294)	5.9% (6; 138)	3.6% (13; 345)
	Long Beach Memorial Medical Center			7.8% (3; 46)	9.7% (12; 132)	1.8% (12; 656)	1.2% (1; 114)	7.4% (27; 381)	3.7% (21; 696)	10.6% (48; 439)	3.1% (11; 397)	1.5% (2; 169)	5.3% (29; 584)
	Los Angeles Community Hospital							5.4% (1; 16)	3.8% (4; 161)	1.7% (1; 95) Better	0% (0; 71)	0% (0; 4)	4.1% (9; 173)
	Los Angeles County/University of Southern California Medical Center	0% (0; 3)	0% (0; 15)		11.1% (17; 250)	3.4% (10; 282)	3.7% (1; 20)	9.9% (15; 213)	3.1% (13; 781)	12.4% (49; 352)	2.4% (7; 361)	0% (0; 53)	3.0% (8; 366)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Los Angeles Metropolitan Medical Center							36.2% (1; 3)	7.5% (11; 165) Worse	5.1% (2; 26)	4.2% (3; 52)	11.4% (1; 4)	5.8% (12; 148)
	Marina Del Rey Hospital						0% (0; 15)	8.3% (5; 36)	2.3% (6; 198)	6.3% (5; 64)	3.1% (4; 102)	2.8% (2; 57)	6.3% (10; 115)
	Memorial Hospital of Gardena				9.5% (3; 11)		0% (0; 3)	7.6% (4; 29)	4.6% (9; 240)	8.4% (14; 107)	2.0% (3; 103)	0% (0; 12)	3.8% (6; 152)
	Methodist Hospital of Southern California			13.6% (1; 9)	19.5% (3; 19)	1.9% (6; 380)	3.4% (1; 29)	7.1% (26; 335)	3.6% (24; 569)	7.1% (25; 326) Better	2.8% (7; 232)	1.8% (3; 201)	4.9% (27; 489)
	Mission Community Hospital – Panorama Campus							5.4% (3; 25)	0% (0; 160) Better	11.9% (5; 37)	2.0% (2; 59)	0% (0; 28)	2.4% (7; 162) Better
	Monterey Park Hospital				0% (0; 4)		0% (0; 3)	9.0% (5; 33)	2.9% (6; 140)	2.0% (1; 50) Better	2.7% (3; 115)	0% (0; 29)	3.9% (3; 100)
	Motion Picture and Television Hospital							33.4% (3; 6) Worse	5.8% (1; 26)	0% (0; 9)	0% (0; 21)		7.2% (2; 44)
	Northridge Hospital Medical Center		0% (0; 3)	0% (0; 13)	6.2% (3; 45)	2.4% (6; 308)	0% (0; 34)	7.4% (18; 252)	1.7% (8; 331) Better	11.9% (29; 229)	0.6% (1; 165)	3.2% (4; 116)	2.8% (11; 292) Better
	Norwalk Community Hospital							14.2% (1; 9)	2.8% (2; 109)	0% (0; 54) Better	4.1% (2; 55)	0% (0; 9)	3.6% (6; 89)
	Olive View – UCLA Medical Center			0% (0; 5)			0% (0; 7)	10.8% (5; 51)	0% (0; 319) Better	2.0% (1; 113) Better	4.7% (4; 141)		2.5% (6; 292)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Olympia Medical Center						0% (0; 3)	3.2% (6; 109) Better	3.8% (15; 327)	5.5% (6; 122) Better	2.3% (6; 160)	2.0% (2; 69)	5.0% (25; 279)
	Pacific Alliance Medical Center, Inc.				0% (0; 5)			2.5% (3; 58) Better	1.2% (5; 262) Better	3.6% (3; 81) Better	0% (0; 73)	3.5% (2; 10)	2.7% (8; 161)
	Pacific Hospital of Long Beach							14.4% (4; 14)	7.0% (7; 87)	10.2% (2; 30)	2.8% (4; 72)	0% (0; 14)	5.6% (11; 115)
	Pacifica Hospital of the Valley							3.6% (1; 13)	5.9% (5; 85)	14.2% (6; 53)	3.4% (2; 64)	9.8% (1; 11)	8.4% (12; 111) Worse
	Pomona Valley Hospital Medical Center			0% (0; 14)	10.5% (10; 83)	1.7% (6; 283)	7.7% (1; 18)	6.9% (23; 369)	2.4% (15; 507)	9.8% (30; 317)	1.6% (7; 285)	6.3% (5; 98)	1.3% (4; 298) Better
	Presbyterian Intercommunity Hospital			0% (0; 12)	2.3% (1; 51)	1.6% (6; 314)	0% (0; 50)	4.3% (20; 425) Better	2.7% (14; 435)	7.2% (18; 256) Better	1.0% (3; 320)	1.5% (2; 151)	2.3% (8; 371) Better
	Providence Holy Cross Medical Center			0% (0; 10)	19.0% (12; 56) Worse	3.0% (7; 327)	0% (0; 63)	11.7% (20; 263) Worse	2.1% (15; 502) Better	11.2% (29; 252)	3.2% (9; 227)	2.7% (3; 104)	2.5% (12; 350) Better
	Providence Saint Joseph Medical Center			0% (0; 9)	6.5% (4; 40)	2.9% (13; 447)	0% (0; 65)	8.2% (20; 293)	2.9% (21; 601)	11.8% (37; 329)	2.0% (8; 262)	0.9% (2; 190)	4.9% (21; 399)
	Providence Tarzana Medical Center			2.4% (1; 43)	13.6% (3; 14)	1.5% (2; 249)	0% (0; 104)	7.7% (10; 158)	3.4% (15; 354)	8.0% (12; 125)	1.9% (3; 150)	6.0% (3; 47)	2.5% (9; 386) Better
	Ronald Reagan UCLA Medical Center	0% (0; 21)	5.2% (2; 56)	1.4% (1; 82)	5.4% (31; 654) Better	3.6% (6; 145)	0% (0; 45)	4.2% (9; 199)	4.5% (23; 448)	8.6% (57; 442) Better	1.1% (1; 123)	0% (0; 38)	4.2% (14; 272)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Saint Francis Medical Center			0% (0; 3)	13.2% (3; 27)	2.2% (3; 137)	0% (0; 18)	8.6% (15; 161)	1.3% (7; 621) Better	7.5% (25; 260) Better	0.9% (3; 259)	0% (0; 43)	3.1% (11; 278)
	Saint John's Health Center		0% (0; 13)	0% (0; 25)	5.6% (3; 61)	1.2% (2; 295)	0% (0; 53)	3.5% (5; 165)	2.5% (12; 300)	8.9% (14; 180)	3.0% (7; 216)	1.8% (2; 129)	4.0% (15; 317)
	Saint Mary Medical Center			0% (0; 6)	19.9% (4; 17)	0% (0; 189)	2.3% (1; 28)	5.9% (8; 168)	4.0% (12; 314)	7.7% (22; 188) Better	2.6% (7; 165)	2.0% (2; 72)	3.2% (16; 345)
	Saint Vincent Medical Center		6.0% (1; 12)	0% (0; 5)	0% (0; 273) Better	1.6% (2; 242)	0% (0; 20)	3.1% (4; 128) Better	1.7% (7; 437)	7.9% (8; 128)	4.0% (4; 119)	2.8% (1; 43)	2.4% (8; 328) Better
	San Dimas Community Hospital (View Comment Letter)						0% (0; 4)	17.9% (9; 32) Worse	1.7% (3; 151)	4.5% (4; 94) Better	0% (0; 99)	1.7% (1; 57)	3.5% (5; 116)
	San Gabriel Valley Medical Center				12.0% (1; 7)	0% (0; 3)	0% (0; 6)	8.2% (16; 91)	1.7% (6; 266)	7.6% (14; 179)	3.1% (6; 154)	2.2% (2; 76)	4.5% (20; 295)
	Santa Monica – UCLA Medical Center and Orthopedic Hospital				27.5% (3; 12)	2.0% (7; 189)	0% (0; 5)	6.7% (11; 158)	1.3% (7; 364) Better	6.0% (9; 150) Better	4.0% (12; 230)	1.5% (3; 124)	4.7% (17; 321)
	Sherman Oaks Hospital							2.8% (5; 68) Better	0.7% (2; 128) Better	9.4% (11; 83)	2.5% (4; 86)	0% (0; 43)	1.2% (2; 96) Better
	Silver Lake Medical Center – Downtown Campus							12.5% (3; 11)	0% (0; 80)	0% (0; 13)	1.7% (1; 67)		1.7% (1; 51)
	Temple Community Hospital							3.7% (2; 38)	2.3% (4; 123)	0% (0; 35)	0% (0; 53)	0% (0; 5)	2.8% (3; 80)
	Torrance Memorial Medical Center (View Comment Letter)		16.8% (1; 3)	0% (0; 17)	11.1% (5; 43)	1.5% (12; 585)	3.0% (2; 80)	8.2% (31; 349)	4.9% (35; 636) Worse	9.7% (35; 349)	1.5% (6; 437)	0.6% (1; 203)	6.1% (30; 489)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Tri-City Regional Medical Center – Hawaiian Gardens							12.3% (4; 14)	2.6% (4; 107)	3.6% (1; 25)	1.8% (1; 64)	5.8% (1; 11)	2.2% (3; 135)
	University of Southern California Kenneth Norris, Jr. Cancer Hospital							0% (0; 4)			3.0% (2; 27)		5.2% (3; 36)
	University of Southern California University Hospital	2.8% (1; 52)	3.2% (1; 38)	2.6% (1; 68)	2.6% (5; 285) Better	1.1% (1; 62)	0% (0; 24)	17.3% (5; 40)	7.2% (11; 89) Worse	4.5% (5; 104) Better	3.1% (1; 40)	0% (0; 7)	5.6% (2; 40)
	Valley Presbyterian Hospital			0% (0; 9)	27.7% (4; 10) Worse	1.7% (8; 242)	9.0% (1; 17)	9.5% (28; 253)	2.5% (9; 329)	13.7% (21; 125)	2.4% (3; 135)	1.1% (1; 82)	7.3% (29; 298) Worse
	Verdugo Hills Hospital				4.1% (1; 16)		20.0% (1; 14)	2.5% (1; 31)	3.7% (6; 175)	8.0% (9; 111)	1.9% (2; 118)	1.4% (1; 92)	8.2% (14; 152) Worse
	West Hills Hospital and Medical Center				32.6% (8; 21) Worse	0.8% (1; 116)	0% (0; 39)	8.5% (10; 130)	3.7% (10; 220)	19.0% (19; 127) Worse	4.9% (10; 185) Worse	2.3% (2; 101)	7.1% (30; 302) Worse
	White Memorial Medical Center				14.1% (3; 15)	1.9% (10; 321)	0% (0; 13)	8.6% (23; 256)	2.2% (11; 467)	6.4% (12; 178) Better	2.8% (9; 238)	0% (0; 58)	3.3% (11; 254)
	Whittier Hospital Medical Center				0% (0; 10)		0% (0; 4)	6.2% (9; 89)	2.8% (7; 231)	6.1% (6; 125)	0.5% (1; 194)	3.5% (2; 82)	4.5% (8; 213)
Madera	Madera Community Hospital							14.5% (17; 85) Worse	9.2% (10; 132) Worse	15.0% (4; 45)	2.0% (1; 64)	0% (0; 12)	18.5% (26; 163) Worse
Marin	Kaiser Foundation Hospital – San Rafael					4.0% (1; 21)	0% (0; 19)	8.8% (12; 102)	4.5% (13; 217)	11.5% (13; 158)	1.9% (2; 138)	3.9% (4; 103)	5.0% (19; 328)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Marin General Hospital			0% (0; 14)	8.7% (4; 29)	0.9% (2; 208)	0% (0; 66)	8.4% (8; 126)	5.2% (17; 221)	14.6% (28; 188) Worse	2.1% (4; 181)	2.4% (3; 102)	5.6% (14; 239)
	Novato Community Hospital							4.0% (1; 14)	1.5% (1; 51)	8.5% (1; 24)	2.3% (1; 32)	0% (0; 24)	1.8% (1; 77)
Mariposa	John C. Fremont Healthcare District								7.0% (1; 19)	11.2% (1; 5)	0% (0; 4)		11.3% (3; 31)
Mendocino	Frank R. Howard Memorial Hospital							17.3% (1; 4)	5.4% (2; 21)	0% (0; 16)	8.1% (1; 15)	6.9% (2; 27)	8.4% (2; 45)
	Mendocino Coast District Hospital							10.6% (1; 5)	10.8% (2; 28)	22.1% (5; 22)	4.9% (1; 17)	0% (0; 4)	2.3% (1; 60)
	Ukiah Valley Medical Center						0% (0; 5)	4.9% (3; 26)	0% (0; 100) Better	11.8% (12; 57)	3.8% (4; 74)	0% (0; 37)	3.6% (5; 98)
Merced	Memorial Hospital Los Banos							8.3% (1; 5)	2.2% (2; 63)	41.3% (2; 17)	0% (0; 32)		9.9% (8; 89)
	Mercy Medical Center – Merced							5.1% (8; 100)	3.4% (21; 404)	6.2% (7; 138)	3.2% (5; 209)	2.9% (3; 94)	4.7% (11; 222)
Modoc	Modoc Medical Center												28.2% (2; 17)
	Surprise Valley Community Hospital												0% (0; 3)
Mono	Mammoth Hospital								0% (0; 5)		0% (0; 5)	0% (0; 5)	0% (0; 8)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
Monterey	Community Hospital Monterey Peninsula	0% (0; 4)	0% (0; 3)	8.6% (1; 6)	10.3% (3; 44)	3.1% (7; 183)	2.4% (1; 37)	9.0% (17; 210)	3.3% (9; 276)	13.9% (24; 181)	3.5% (8; 226)	2.1% (2; 116)	5.7% (18; 299)
	George L. Mee Memorial Hospital							6.5% (1; 9)	5.8% (2; 46)		0% (0; 15)		3.9% (2; 64)
	Natividad Medical Center							8.8% (4; 22)	0% (0; 86)	0% (0; 35)	0% (0; 83)	0% (0; 5)	5.0% (4; 133)
	Salinas Valley Memorial Hospital	9.2% (1; 4)	0% (0; 3)	0% (0; 8)	6.3% (3; 39)	1.5% (4; 260)	0% (0; 29)	6.4% (12; 203)	2.2% (7; 315)	8.7% (16; 180)	3.9% (9; 211)	7.9% (8; 103) Worse	4.3% (12; 238)
Napa	N.M. Holderman Memorial Hospital(Veteran's Home of California) (View Comment Letter)								0% (0; 4)				0% (0; 16)
	Queen of the Valley Hospital – Napa			0% (0; 19)		2.7% (9; 238)	0% (0; 36)	7.1% (16; 220)	2.1% (6; 200)	11.0% (12; 111)	1.8% (3; 160)	2.8% (3; 89)	4.8% (13; 245)
	Saint Helena Hospital			0% (0; 13)		3.1% (4; 379)	0% (0; 66)	9.0% (5; 69)	3.7% (3; 94)	13.5% (6; 38)	1.2% (1; 66)	5.0% (1; 24)	6.1% (4; 80)
Nevada	Sierra Nevada Memorial Hospital			0% (0; 3)			0% (0; 17)	9.0% (6; 37)	5.1% (14; 179)	11.3% (14; 99)	1.8% (3; 126)	1.1% (1; 86)	4.0% (9; 236)
	Tahoe Forest Hospital						0% (0; 9)	0% (0; 6)	3.0% (1; 40)	32.8% (3; 10)	0% (0; 26)	0% (0; 25)	3.6% (1; 44)
Orange	Anaheim General Hospital							7.5% (2; 14)	4.4% (1; 25)	0% (0; 13)	2.0% (1; 72)	12.4% (2; 9)	1.2% (3; 162) Better
	Anaheim Memorial Medical Center**			6.9% (1; 21)	13.1% (3; 23)	1.6% (6; 510)	0% (0; 56)	6.7% (19; 362)	2.9% (15; 510)	7.5% (16; 249)	1.1% (2; 229)	1.6% (2; 97)	2.9% (12; 387)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Chapman Medical Center							14.1% (4; 6)	2.9% (2; 47)	0% (0; 13)	4.5% (2; 31)	0% (0; 9)	3.4% (4; 101)
	Coastal Communities Hospital						0% (0; 3)	5.8% (2; 17)	1.8% (1; 61)	16.2% (3; 21)	2.4% (3; 73)	0% (0; 6)	3.2% (3; 105)
	College Hospital Costa Mesa												0% (0; 5)
	Fountain Valley Regional Hospital and Medical Center – Euclid	0% (0; 6)	0% (0; 3)	0% (0; 17)	8.5% (7; 81)	2.3% (8; 260)	0% (0; 19)	6.7% (16; 264)	2.5% (11; 405)	9.9% (29; 251)	2.7% (6; 271)	2.8% (2; 67)	2.9% (11; 332)
	Garden Grove Hospital and Medical Center				10.0% (3; 9)		0% (0; 9)	8.8% (9; 54)	0% (0; 109) Better	5.6% (7; 114) Better	1.8% (2; 115)	3.9% (2; 45)	3.5% (8; 215)
	Hoag Memorial Hospital Presbyterian	0% (0; 4)	0% (0; 3)	0% (0; 24)	7.5% (20; 300)	1.8% (6; 556)	0% (0; 99)	6.7% (17; 280)	3.4% (25; 575)	9.2% (51; 468)	2.7% (10; 340)	2.1% (5; 256)	6.7% (44; 562) Worse
	Huntington Beach Hospital							6.3% (4; 26)	1.4% (2; 97)	8.5% (3; 44)	3.5% (3; 61)	1.3% (1; 21)	3.3% (5; 112)
	Irvine Medical Center				0% (0; 9)	1.4% (2; 240)		5.0% (6; 181)	2.8% (5; 154)	9.6% (3; 85)	3.9% (4; 119)	5.6% (2; 41)	4.5% (8; 167)
	Kaiser Foundation Hospital – Anaheim		0% (0; 15)	0% (0; 20)	8.4% (8; 157)		0% (0; 40)	5.0% (10; 135)	2.7% (14; 378)	7.2% (28; 388) Better	0% (0; 284) Better	1.6% (3; 156)	3.9% (17; 337)
	La Palma Intercommunity Hospital (View Comment Letter)						0% (0; 5)	1.1% (1; 70) Better	2.2% (2; 98)	3.9% (2; 78)	0% (0; 57)	0% (0; 31)	2.0% (1; 47)
	Los Alamitos Medical Center			8.0% (1; 11)	8.8% (5; 30)	2.3% (2; 80)	0% (0; 66)	2.5% (3; 154) Better	3.6% (13; 343)	9.6% (22; 213)	2.7% (3; 189)	3.3% (5; 145)	5.0% (13; 284)
	Mission Hospital Laguna Beach				0% (0; 27)		0% (0; 13)	10.6% (4; 14)	1.3% (1; 71)	10.0% (5; 34)	2.8% (1; 49)	0% (0; 37)	3.6% (3; 68)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Mission Hospital Regional Medical Center	0% (0; 3)		0% (0; 8)	5.5% (7; 115)	2.3% (9; 560)	0% (0; 18)	6.4% (16; 298)	2.7% (12; 349)	9.5% (34; 314)	2.8% (7; 229)	2.2% (3; 136)	5.0% (14; 305)
	Orange Coast Memorial Medical Center (View Comment Letter)				20.4% (5; 18) Worse	2.1% (1; 23)	0% (0; 34)	5.9% (14; 132)	3.1% (14; 369)	6.8% (16; 213) Better	2.4% (5; 218)	2.7% (3; 146)	5.4% (16; 301)
	Placentia Linda Hospital							12.2% (8; 39)	4.2% (6; 141)	12.1% (4; 65)	4.3% (3; 100)	1.3% (1; 63)	4.2% (8; 222)
	Saddleback Memorial Medical Center	0% (0; 3)		0% (0; 23)	4.1% (3; 51)	2.2% (10; 320)	0% (0; 37)	6.1% (30; 325)	2.5% (16; 457)	8.8% (27; 284)	0.9% (3; 259)	0.8% (2; 234) Better	3.2% (14; 351)
	Saddleback Memorial Medical Center – San Clemente							8.8% (3; 16)	1.3% (2; 113)	5.8% (4; 64)	3.0% (2; 67)	0% (0; 54)	4.1% (6; 138)
	Saint Joseph Hospital – Orange		0% (0; 4)	0% (0; 44)	4.0% (6; 127)	2.8% (8; 258)	0% (0; 110)	7.2% (24; 294)	3.1% (18; 481)	7.9% (26; 338)	2.1% (8; 304)	0.7% (1; 149)	4.3% (18; 371)
	Saint Jude Medical Center			0% (0; 21)	5.9% (3; 59)	2.3% (10; 559)	0% (0; 60)	8.4% (31; 359)	3.2% (17; 446)	10.7% (34; 305)	2.0% (6; 292)	2.8% (7; 186)	2.5% (10; 344) Better
	UC Irvine Medical Center	0% (0; 15)	15.3% (2; 17)	0% (0; 20)	7.8% (23; 207)	2.5% (5; 172)	0% (0; 21)	10.1% (10; 102)	2.2% (5; 228)	9.0% (53; 275)	1.1% (2; 133)	0% (0; 28)	3.1% (7; 163)
	West Anaheim Medical Center					1.5% (3; 120)	0% (0; 6)	5.8% (10; 148)	2.4% (11; 262)	4.7% (5; 107) Better	0.7% (1; 132)	1.1% (1; 63)	1.4% (4; 202) Better
	Western Medical Center – Anaheim					1.1% (2; 172)	3.1% (1; 13)	7.7% (12; 188)	6.0% (4; 72)	12.3% (2; 28)	3.4% (1; 33)	0% (0; 10)	1.5% (1; 82)
	Western Medical Center – Santa Ana				12.5% (10; 78)	1.4% (4; 201)	0% (0; 9)	10.3% (18; 194)	5.0% (8; 189)	11.8% (40; 209)	2.5% (3; 75)	7.3% (4; 42)	3.2% (5; 177)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
Placer	Sutter Auburn Faith Hospital						0% (0; 7)	0% (0; 10)	5.3% (10; 125)	11.6% (7; 84)	0% (0; 67)	2.7% (3; 82)	1.5% (2; 149)
	Sutter Roseville Medical Center			0% (0; 21)	9.0% (9; 71)	2.3% (3; 137)	0% (0; 46)	6.0% (17; 226)	2.4% (7; 260)	10.9% (39; 305)	2.8% (7; 257)	3.2% (8; 203)	5.0% (13; 280)
Plumas	Eastern Plumas Hospital – Portola Campus							0% (0; 4)	4.7% (1; 21)	80.3% (1; 3)	0% (0; 5)		7.0% (1; 22)
	Plumas District Hospital								5.6% (1; 24)		0% (0; 18)		0% (0; 41)
	Seneca Healthcare District								6.8% (1; 13)				8.9% (1; 29)
Riverside	Corona Regional Medical Center – Main			0% (0; 3)			0% (0; 10)	11.0% (10; 57)	1.2% (3; 220) Better	9.7% (10; 111)	1.8% (3; 134)	2.7% (2; 94)	5.2% (17; 282)
	Desert Regional Medical Center			0% (0; 24)	10.8% (6; 54)	1.6% (9; 581)	0% (0; 58)	6.1% (17; 357)	3.9% (14; 348)	10.6% (50; 350)	4.5% (12; 238) Worse	4.2% (5; 139)	6.3% (15; 290)
	Eisenhower Medical Center			0% (0; 36)	5.6% (4; 69)	2.3% (13; 635)	0% (0; 153)	7.8% (36; 500)	2.6% (18; 732)	7.4% (32; 496) Better	2.9% (12; 404)	2.2% (6; 251)	3.9% (21; 590)
	Hemet Valley Medical Center (View Comment Letter)			0% (0; 10)			3.4% (2; 61)	9.0% (32; 202)	5.7% (23; 431) Worse	12.2% (26; 227)	3.6% (10; 251)	2.8% (6; 188)	5.9% (23; 340)
	John F. Kennedy Memorial Hospital					3.6% (1; 25)	0% (0; 14)	6.0% (6; 92)	0% (0; 207) Better	8.7% (4; 69)	0% (0; 86)	2.1% (1; 66)	5.4% (10; 238)
	Kaiser Foundation Hospital – Moreno Valley						0% (0; 7)	10.0% (6; 16)	0.8% (1; 103)	8.8% (3; 41)	3.1% (2; 61)	13.4% (1; 12)	2.9% (6; 161)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Kaiser Foundation Hospital – Riverside	14.2% (1; 11)	0% (0; 6)	0% (0; 10)			0% (0; 44)	7.5% (14; 145)	1.7% (8; 380) Better	5.8% (10; 209) Better	2.0% (4; 263)	2.4% (2; 110)	3.5% (15; 417)
	Menifee Valley Medical Center (View Comment Letter)						0% (0; 21)	11.8% (28; 125) Worse	3.4% (6; 175)	4.9% (3; 78)	4.4% (3; 93)	4.0% (2; 84)	5.4% (9; 188)
	Palo Verde Hospital							0% (0; 3)	6.9% (1; 30)	0% (0; 8)	0% (0; 21)		15.8% (4; 58)
	Parkview Community Hospital Medical Center						0% (0; 7)	7.5% (7; 53)	1.1% (3; 181) Better	4.7% (3; 102)	2.2% (3; 113)	1.0% (1; 66)	2.1% (11; 308) Better
	Riverside Community Hospital			0% (0; 21)	4.6% (2; 66)	2.6% (22; 830)	1.4% (1; 66)	10.0% (46; 588) Worse	3.2% (23; 658)	9.0% (38; 289)	1.7% (6; 335)	2.1% (3; 135)	6.7% (23; 364)
	Riverside County Regional Medical Center				7.6% (3; 55)		0% (0; 17)	11.0% (5; 52)	1.3% (3; 415)	5.4% (11; 194) Better	0% (0; 124)	3.9% (2; 34)	5.1% (11; 293)
	San Geronio Memorial Hospital							5.8% (10; 88)	2.2% (6; 198)	8.1% (5; 85)	3.4% (6; 104)	0% (0; 42)	4.9% (12; 253)
	Southwest Healthcare System – Murrieta			0% (0; 24)	6.0% (5; 67)		0% (0; 66)	6.8% (12; 111)	1.9% (9; 460)	8.0% (22; 290)	2.0% (4; 262)	1.3% (2; 156)	5.1% (16; 416)
Sacramento	Kaiser Foundation Hospital – Sacramento/Roseville-Morse	10.0% (2; 15)	0% (0; 12)	9.7% (2; 45)	5.0% (11; 267)		1.8% (2; 155)	7.9% (35; 333)	4.9% (48; 988) Worse	11.7% (89; 782)	3.1% (17; 532)	5.1% (14; 348) Worse	7.7% (72; 902) Worse
	Kaiser Foundation Hospital – South Sacramento	0% (0; 3)	0% (0; 8)	11.9% (2; 27)	0% (0; 3)		0% (0; 31)	8.1% (16; 172)	2.9% (10; 382)	8.1% (17; 235)	7.9% (8; 173) Worse	5.4% (5; 120)	6.6% (19; 329)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Mercy General Hospital			0% (0; 19)	3.0% (1; 43)	2.1% (24; 1,647)	1.9% (3; 79)	5.7% (32; 1,090)	2.7% (13; 514)	7.1% (16; 253) Better	1.3% (4; 194)	0.9% (1; 112)	3.2% (10; 308)
	Mercy Hospital – Folsom							6.5% (3; 18)	2.0% (4; 131)	7.4% (7; 83)	2.6% (4; 101)	7.4% (3; 62)	2.0% (3; 118)
	Mercy San Juan Hospital			0% (0; 26)	7.9% (6; 54)	1.7% (4; 250)	2.7% (2; 69)	4.7% (17; 322) Better	2.1% (14; 463) Better	11.8% (59; 412)	1.8% (8; 323)	1.6% (4; 207)	2.1% (9; 402) Better
	Methodist Hospital of Sacramento						0% (0; 6)	7.3% (4; 28)	4.0% (12; 250)	8.7% (18; 180)	1.1% (2; 195)	2.3% (2; 70)	6.2% (17; 243)
	Sutter General Hospital	6.0% (1; 14)	0% (0; 6)	0% (0; 7)	3.9% (5; 111)		0% (0; 33)	8.5% (7; 31)	1.1% (3; 237) Better	8.2% (29; 292)	2.4% (8; 276)	0.9% (2; 171)	3.3% (13; 417)
	Sutter Memorial Hospital			0% (0; 41)	0% (0; 89)	2.4% (11; 448)	1.8% (2; 98)	8.0% (24; 413)	3.4% (11; 333)	12.7% (11; 74)	3.2% (5; 79)		1.9% (3; 166)
	UC Davis Medical Center	0% (0; 8)	4.4% (1; 23)	9.0% (2; 26)	7.5% (10; 280)	1.7% (10; 483)	0% (0; 47)	5.6% (19; 410)	2.1% (11; 621)	12.2% (46; 324)	4.5% (14; 319) Worse	3.6% (3; 59)	2.5% (11; 443) Better
San Benito	Hazel Hawkins Memorial Hospital							0% (0; 12)	3.4% (4; 99)	7.0% (2; 25)	0% (0; 34)	0% (0; 20)	7.2% (5; 79)
San Bernardino	Arrowhead Regional Medical Center	0% (0; 3)			15.1% (14; 96) Worse	0% (0; 88)	0% (0; 14)	9.4% (9; 137)	2.8% (6; 397)	13.9% (60; 341) Worse	1.5% (2; 192)	3.6% (1; 50)	7.8% (20; 343) Worse
	Barstow Community Hospital							13.4% (8; 44)	3.1% (5; 194)	3.0% (1; 68)	3.7% (2; 60)	0% (0; 3)	0% (0; 120)
	Bear Valley Community Hospital							0% (0; 6)	0% (0; 24)	0% (0; 10)	0% (0; 11)		0% (0; 30)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Chino Valley Medical Center							5.0% (4; 31)	0.4% (1; 190) Better	0% (0; 114) Better	0.8% (1; 88)	0% (0; 32)	0.9% (2; 148) Better
	Colorado River Medical Center							0% (0; 3)	0% (0; 19)	63.9% (1; 7)	11.1% (1; 18)		0% (0; 11)
	Community Hospital of San Bernardino						13.8% (1; 9)	3.2% (2; 46)	1.5% (2; 189)	4.1% (2; 96)	0.9% (1; 101)	2.2% (1; 17)	2.2% (6; 238) Better
	Desert Valley Hospital							3.5% (3; 36)	0% (0; 218) Better	2.2% (3; 121) Better	0.6% (1; 130)	0% (0; 48) Better	0.7% (2; 225) Better
	Hi-Desert Medical Center						0% (0; 6)	0% (0; 7)	2.0% (2; 104)	15.0% (6; 79)	0% (0; 47)	1.5% (1; 58)	3.4% (5; 194)
	Kaiser Foundation Hospital – Fontana	0% (0; 4)	11.2% (1; 7)	0% (0; 21)	7.3% (6; 95)		0% (0; 50)	7.2% (21; 216)	2.3% (20; 692)	11.0% (56; 528)	2.5% (11; 517)	4.4% (10; 184)	3.5% (36; 737) Better
	Loma Linda University Medical Center	70.0% (1; 4)	0% (0; 7)	6.0% (1; 39)	8.5% (15; 289)	3.1% (7; 258)	0% (0; 23)	6.8% (21; 397)	3.0% (14; 502)	8.4% (39; 362) Better	0.5% (1; 174)	4.8% (5; 90)	5.5% (19; 366)
	Montclair Hospital Medical Center							6.5% (2; 19)	3.1% (5; 121)	3.4% (1; 44)	0% (0; 59)	2.8% (1; 25)	1.6% (2; 94)
	Mountains Community Hospital								0% (0; 12)		0% (0; 9)		0% (0; 27)
	Redlands Community Hospital	8.5% (1; 5)		0% (0; 4)			7.8% (2; 54) Worse	8.4% (15; 124)	3.3% (10; 247)	4.8% (6; 129) Better	1.6% (3; 183)	4.3% (4; 136)	4.9% (27; 449)
	Saint Bernadine Medical Center			0% (0; 21)	0% (0; 29)	1.9% (17; 1,630)	0% (0; 41)	6.7% (41; 970)	3.7% (20; 592)	6.0% (13; 205) Better	1.5% (3; 180)	1.0% (1; 103)	3.4% (13; 344)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Saint Mary Regional Medical Center			0% (0; 12)	0% (0; 10)	1.1% (3; 308)	2.7% (1; 59)	7.1% (21; 396)	1.5% (8; 588) Better	2.5% (2; 207) Better	1.2% (2; 208)	3.8% (5; 115)	6.7% (21; 365)
	San Antonio Community Hospital			0% (0; 13)	10.0% (12; 61)	2.0% (8; 265)	1.0% (1; 52)	7.4% (23; 308)	1.6% (10; 488) Better	7.0% (26; 297) Better	2.2% (6; 243)	2.0% (3; 132)	2.8% (13; 478) Better
	Victor Valley Community Hospital						0% (0; 5)	2.8% (1; 40)	3.5% (5; 150)	0% (0; 43) Better	0% (0; 63)	8.2% (2; 37)	3.6% (4; 123)
San Diego	Alvarado Hospital			0% (0; 8)	11.4% (5; 40)	1.0% (4; 445)	0% (0; 39)	6.4% (8; 138)	1.4% (4; 308)	10.4% (21; 147)	1.8% (4; 165)	2.1% (1; 73)	4.3% (15; 359)
	Fallbrook Hospital District							9.1% (2; 12)	3.0% (2; 62)	14.5% (2; 21)	1.4% (1; 52)	0% (0; 34)	1.7% (1; 76)
	Grossmont Hospital			8.4% (2; 25)	7.9% (10; 110)	1.7% (15; 682)	0% (0; 105)	8.3% (48; 533)	4.7% (31; 688)	11.4% (73; 550)	2.7% (10; 402)	4.8% (12; 266)	4.9% (28; 699)
	Kaiser Foundation Hospital – San Diego	0% (0; 7)	39.4% (1; 5)	4.4% (1; 28)	4.8% (5; 190)		0% (0; 62)	7.4% (18; 165)	2.3% (21; 819)	10.1% (40; 482)	1.3% (4; 432)	2.5% (8; 315)	5.9% (52; 953)
	Palomar Medical Center	111.3% (1; 3)		0% (0; 22)	13.6% (6; 39)	1.5% (6; 427)	0% (0; 33)	7.6% (24; 401)	3.5% (17; 515)	11.9% (53; 446)	2.6% (6; 305)	4.2% (7; 194)	3.9% (13; 435)
	Paradise Valley Hospital (View Comment Letter)			0% (0; 6)	18.0% (2; 5)		5.3% (1; 20)	6.9% (9; 63)	1.4% (5; 303) Better	5.0% (8; 124) Better	3.2% (3; 81)	0% (0; 25)	2.8% (6; 203)
	Pomerado Hospital			0% (0; 6)	43.4% (4; 11) Worse		0% (0; 6)	12.2% (11; 57)	6.2% (13; 213)	14.5% (19; 145)	2.7% (4; 180)	5.0% (5; 121)	8.7% (19; 239) Worse
	Scripps Green Hospital		0% (0; 3)	13.1% (2; 37)	5.8% (1; 55)	1.9% (12; 1,035)	6.3% (1; 53)	6.9% (10; 190)	3.5% (9; 307)	7.5% (6; 120)	0% (0; 175)	3.2% (4; 108)	3.7% (8; 229)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Scripps Memorial Hospital – Encinitas			0% (0; 12)	8.6% (3; 28)	1.2% (3; 133)	0% (0; 27)	4.7% (8; 165)	2.8% (6; 223)	7.4% (11; 153)	1.5% (2; 152)	1.8% (2; 89)	2.6% (6; 202)
	Scripps Memorial Hospital – La Jolla		26.1% (1; 5)	0% (0; 8)	9.1% (7; 103)	2.2% (14; 1,156)	0% (0; 25)	9.9% (30; 496)	2.6% (9; 283)	10.7% (31; 241)	3.2% (4; 152)	2.1% (4; 126)	2.6% (6; 227)
	Scripps Mercy Hospital			0% (0; 6)	10.9% (11; 85)	0.2% (1; 403) Better	0% (0; 40)	4.7% (11; 263)	3.2% (15; 432)	9.4% (41; 314)	2.2% (5; 228)	2.0% (3; 145)	3.0% (9; 297)
	Scripps Mercy Hospital – Chula Vista			0% (0; 4)	11.1% (1; 9)	0.8% (1; 105)	0% (0; 6)	7.0% (15; 147)	1.9% (11; 421) Better	8.0% (14; 194)	2.1% (5; 199)	4.1% (5; 66)	2.4% (8; 302) Better
	Sharp Chula Vista Medical Center			0% (0; 12)	9.3% (7; 50)	0.9% (5; 424) Better	1.5% (1; 53)	5.6% (21; 313)	1.8% (13; 566) Better	9.0% (25; 292)	1.9% (5; 230)	0.9% (1; 88)	3.5% (16; 374)
	Sharp Coronado Hospital and Healthcare Center							14.3% (3; 8)	1.0% (1; 61)	24.0% (2; 11)	2.6% (1; 34)	6.5% (2; 29)	6.4% (8; 92)
	Sharp Memorial Hospital (View Comment Letter)	0% (0; 4)	0% (0; 3)	5.3% (1; 31)	10.1% (8; 76)	2.7% (8; 346)	1.8% (1; 50)	9.5% (32; 365)	4.1% (21; 513)	8.7% (27; 294)	2.6% (7; 285)	3.3% (5; 183)	5.1% (26; 509)
	Tri-City Medical Center – Oceanside			0% (0; 21)	12.9% (8; 58)	3.0% (4; 312)	3.6% (2; 71)	10.4% (34; 355)	4.8% (20; 419)	14.2% (63; 362) Worse	4.5% (13; 301) Worse	2.1% (4; 260)	6.6% (38; 542) Worse
	UC San Diego Medical Center	21.1% (1; 7)	0% (0; 31)	12.8% (2; 17)	6.7% (16; 251)	0.8% (3; 325)	2.7% (1; 35)	3.7% (8; 234) Better	2.7% (12; 385)	7.1% (31; 314) Better	0.8% (2; 234)	1.4% (2; 88)	3.1% (11; 350)
San Francisco	California Pacific Medical Center – Pacific Campus	0% (0; 6)	3.4% (1; 26)	4.3% (1; 28)	7.2% (14; 190)	2.2% (9; 395)	0% (0; 43)	8.7% (18; 253)	4.1% (26; 562)	8.3% (38; 382)	3.7% (14; 334)	1.5% (2; 158)	4.0% (20; 464)
	California Pacific Medical Center – St. Luke’s Campus						0% (0; 7)	10.3% (3; 14)	5.4% (9; 189)	9.9% (5; 48)	0% (0; 68)	2.6% (1; 37)	5.1% (6; 131)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Chinese Hospital							10.5% (3; 20)	0% (0; 121) Better	5.5% (2; 70)	0% (0; 110)	0% (0; 18)	4.7% (9; 235)
	Kaiser Foundation Hospital – San Francisco	0% (0; 3)	42.7% (1; 5)	0% (0; 57)		1.2% (11; 1,209)	0% (0; 74)	5.0% (22; 815) Better	2.6% (12; 430)	13.9% (25; 232)	1.2% (2; 185)	2.1% (1; 58)	4.2% (13; 250)
	Laguna Honda Hospital and Rehabilitation Center								0% (0; 4)				8.8% (4; 36)
	Saint Francis Memorial Hospital				7.1% (2; 21)		0% (0; 7)	12.2% (4; 25)	4.8% (11; 205)	10.2% (13; 114)	4.6% (3; 80)	0% (0; 59)	10.4% (21; 193) Worse
	Saint Mary’s Medical Center, San Francisco			0% (0; 7)	29.0% (4; 9) Worse	3.4% (11; 319) Worse	0% (0; 9)	9.8% (13; 167)	3.8% (11; 216)	8.9% (10; 98)	3.6% (8; 148)	0% (0; 68)	3.3% (11; 227)
	San Francisco General Hospital				17.0% (11; 49) Worse	6.0% (3; 58)		10.0% (10; 115)	4.2% (7; 311)	12.6% (32; 164)	2.3% (3; 174)	0% (0; 37)	3.2% (8; 424)
	UC San Francisco Medical Center	2.4% (1; 36)	0% (0; 22)	9.9% (3; 41) Worse	7.9% (55; 1,209)	2.0% (5; 188)	2.9% (1; 55)	6.2% (16; 184)	2.6% (15; 441)	13.4% (82; 484) Worse	1.3% (3; 194)	3.6% (4; 77)	3.1% (14; 429)
San Joaquin	Dameron Hospital					1.8% (1; 165)	0% (0; 51)	14.2% (20; 217) Worse	6.0% (19; 308) Worse	13.1% (18; 133)	0.8% (1; 151)	2.3% (2; 105)	7.9% (33; 316) Worse
	Doctors Hospital of Manteca							11.6% (3; 19)	4.0% (7; 172)	16.0% (8; 57)	0% (0; 90)	6.4% (3; 38)	12.4% (9; 108) Worse
	Kaiser Foundation Hospital – Manteca						0% (0; 6)	5.4% (2; 38)	3.8% (6; 189)	10.7% (6; 75)	2.9% (3; 115)	5.1% (3; 42)	8.6% (12; 156)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Lodi Memorial Hospital			0% (0; 4)			0% (0; 26)	11.9% (10; 51)	4.6% (11; 232)	12.0% (17; 120)	5.8% (4; 125)	1.4% (1; 95)	6.1% (14; 259)
	Saint Joseph's Medical Center of Stockton	0% (0; 5)		9.2% (2; 39)	10.0% (7; 60)	1.7% (5; 584)	0% (0; 129)	9.8% (24; 307)	2.4% (14; 561)	10.9% (41; 266)	1.7% (5; 285)	1.7% (2; 137)	7.2% (40; 554) Worse
	San Joaquin General Hospital				22.4% (2; 11)			9.1% (6; 74)	1.7% (3; 253)	12.4% (21; 133)	3.6% (6; 168)	7.6% (1; 25)	4.3% (6; 202)
	Sutter Tracy Community Hospital							12.7% (3; 11)	3.9% (4; 112)	18.4% (10; 67)	1.1% (1; 86)	4.4% (1; 38)	4.4% (4; 145)
San Luis Obispo	Arroyo Grande Community Hospital						0% (0; 20)	3.0% (1; 22)	7.0% (7; 103)	13.2% (10; 51)	5.2% (4; 81)	7.0% (4; 68)	5.7% (8; 126)
	French Hospital Medical Center					1.8% (3; 305)	0% (0; 34)	6.9% (4; 155)	4.0% (5; 97)	11.3% (4; 69)	1.7% (2; 126)	0% (0; 45)	4.8% (6; 146)
	Sierra Vista Regional Medical Center (View Comment Letter)			0% (0; 11)	8.9% (5; 90)	3.7% (4; 112)	0% (0; 23)	9.7% (7; 98)	2.2% (2; 93)	17.2% (28; 161) Worse	7.8% (4; 85)	4.7% (2; 58)	3.2% (3; 146)
	Twin Cities Community Hospital						0% (0; 13)	10.6% (6; 33)	5.7% (8; 164)	15.7% (12; 117)	1.2% (1; 135)	3.3% (2; 81)	5.3% (6; 206)
San Mateo	Kaiser Foundation Hospital – Redwood City (View Comment Letter)				7.9% (36; 645)			4.7% (6; 65)	6.1% (17; 164) Worse	13.0% (82; 458) Worse	2.6% (3; 111)	5.6% (4; 68)	3.8% (12; 242)
	Kaiser Foundation Hospital – South San Francisco						0% (0; 16)	6.9% (8; 64)	2.9% (10; 222)	12.1% (18; 154)	2.5% (5; 162)	2.5% (3; 94)	4.3% (21; 319)
	Peninsula Medical Center	0% (0; 3)	17.1% (1; 6)	10.1% (1; 17)	4.0% (1; 17)	2.2% (7; 238)	0% (0; 42)	4.8% (10; 209)	2.7% (13; 351)	8.1% (28; 267) Better	1.8% (7; 263)	4.2% (7; 163)	2.7% (9; 310)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	San Mateo Medical Center				0% (0; 11)		0% (0; 7)	8.3% (2; 31)	2.3% (2; 130)	9.3% (4; 36)	0% (0; 65)	0% (0; 11)	6.1% (6; 134)
	Sequoia Hospital			0% (0; 15)	0% (0; 8)	1.4% (2; 208)	0% (0; 42)	7.3% (8; 111)	5.2% (17; 247)	12.8% (11; 75)	1.2% (1; 83)	10.3% (6; 75) Worse	3.0% (4; 127)
	Seton Medical Center			15.6% (2; 5) Worse	10.4% (5; 17)	1.8% (8; 572)	0% (0; 26)	11.4% (17; 156)	3.0% (18; 530)	10.3% (32; 189)	3.4% (6; 193)	2.4% (2; 82)	2.5% (12; 363) Better
Santa Barbara	Goleta Valley Cottage Hospital								2.8% (1; 30)	35.3% (2; 9)	0% (0; 4)	0% (0; 78)	4.9% (2; 54)
	Lompoc Valley Medical Center						15.7% (1; 14)	0% (0; 7)	0% (0; 50)	18.8% (10; 49) Worse	2.3% (1; 56)	4.0% (1; 35)	15.9% (13; 90) Worse
	Marian Medical Center			0% (0; 5)	16.5% (2; 21)	1.5% (1; 125)	7.1% (2; 54) Worse	9.2% (12; 190)	7.5% (16; 212) Worse	23.1% (33; 190) Worse	2.2% (6; 214)	4.9% (6; 123)	5.6% (9; 241)
	Santa Barbara Cottage Hospital	22.8% (1; 11)	0% (0; 12)	12.4% (1; 23)	10.2% (18; 178)	3.7% (12; 467) Worse	0% (0; 65)	11.7% (19; 220)	4.2% (18; 369)	12.5% (55; 424)	2.9% (7; 267)	3.7% (5; 175)	4.4% (15; 386)
	Santa Ynez Valley Cottage Hospital								3.5% (1; 18)	18.2% (1; 8)	0% (0; 7)		4.6% (1; 33)
Santa Clara	Community Hospital of Los Gatos**				0% (0; 3)			21.1% (8; 14) Worse	2.1% (2; 90)	11.4% (1; 19)	2.8% (1; 54)	2.3% (1; 45)	8.0% (8; 103)
	El Camino Hospital	0% (0; 3)		5.8% (1; 39)	5.2% (2; 36)	2.7% (10; 271)	0% (0; 38)	9.3% (21; 242)	4.8% (24; 430)	17.1% (40; 242) Worse	1.6% (4; 337)	2.2% (3; 173)	6.0% (19; 357)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Good Samaritan Hospital – San Jose	17.2% (1; 3)	9.0% (1; 8)	0% (0; 4)	12.1% (15; 99) Worse	2.5% (16; 602)	0% (0; 44)	7.7% (17; 361)	5.0% (21; 307)	11.7% (33; 301)	3.5% (7; 190)	3.6% (5; 138)	7.1% (19; 298)
	Kaiser Foundation Hospital – San Jose		42.1% (1; 6)	0% (0; 6)		3.8% (4; 60)	0% (0; 38)	10.9% (23; 181)	4.1% (21; 378)	14.7% (28; 234)	1.7% (3; 213)	0.9% (1; 97)	5.2% (22; 370)
	Kaiser Foundation Hospital – Santa Clara	15.8% (1; 7)	0% (0; 3)	4.7% (1; 46)		1.8% (14; 982)	0% (0; 71)	6.2% (24; 651)	3.6% (29; 627)	14.3% (38; 298) Worse	1.9% (6; 299)	0.6% (1; 153)	6.4% (36; 428)
	O’Connor Hospital – San Jose			11.7% (1; 15)	0% (0; 32)	3.9% (9; 309) Worse	0% (0; 40)	10.4% (11; 141)	3.2% (14; 407)	7.1% (13; 171)	2.1% (4; 220)	2.6% (2; 92)	8.8% (44; 383) Worse
	Regional Medical of San Jose			0% (0; 3)	11.4% (9; 38)	3.8% (9; 231) Worse		12.2% (18; 156) Worse	3.7% (14; 408)	11.1% (42; 277)	3.0% (7; 238)	2.6% (2; 64)	5.1% (16; 345)
	Saint Louise Regional Hospital							6.6% (2; 27)	4.1% (4; 105)	10.0% (3; 34)	2.7% (1; 50)	3.1% (1; 37)	5.8% (7; 131)
	Santa Clara Valley Medical Center		22.4% (1; 5)	0% (0; 3)	15.1% (13; 86) Worse	2.1% (7; 224)	0% (0; 16)	9.5% (15; 179)	2.9% (10; 483)	9.7% (25; 284)	0% (0; 255) Better	2.8% (2; 49)	2.9% (11; 359)
	Stanford Hospital	0% (0; 15)	3.5% (1; 32)	0% (0; 85)	5.3% (37; 788) Better	1.7% (6; 448)	0% (0; 84)	7.2% (13; 202)	5.1% (30; 420) Worse	9.0% (49; 428)	3.4% (9; 246)	1.5% (2; 143)	4.2% (16; 392)
Santa Cruz	Dominican Hospital – Santa Cruz/Soquel		0% (0; 3)	0% (0; 5)	9.4% (5; 56)	3.3% (6; 310)	0% (0; 31)	9.0% (18; 234)	3.8% (13; 324)	13.4% (31; 228)	5.6% (14; 210) Worse	6.1% (6; 117)	4.3% (11; 312)
	Watsonville Community Hospital							7.0% (6; 58)	6.0% (8; 194)	15.1% (7; 60)	3.3% (3; 98)	8.9% (5; 49) Worse	8.0% (9; 170)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
Shasta	Mayers Memorial Hospital								31.9% (4; 22) Worse	123.5% (1; 3)	23.9% (1; 7)	7.6% (1; 8)	5.6% (1; 27)
	Mercy Medical Center – Redding	21.6% (2; 3)		11.8% (2; 32)	9.8% (6; 79)	1.9 (6; 373)	0% (0; 124)	8.2% (24; 309)	3.4% (16; 322)	8.6% (30; 276)	1.7% (4; 159)	0.6% (1; 142)	2.9% (11; 369)
	Shasta Regional Medical Center (View Comment Letter)			0% (0; 6)	74.4% (1; 6)	4.2% (11; 187) Worse	0% (0; 50)	8.0% (11; 179)	4.0% (13; 285)	8.5% (11; 162)	2.6% (5; 192)	6.8% (5; 94)	9.9% (16; 258) Worse
Siskiyou	Fairchild Medical Center						0% (0; 6)	11.0% (2; 8)	2.8% (1; 36)	5.1% (1; 27)	0% (0; 36)	2.7% (1; 33)	4.1% (4; 135)
	Mercy Medical Center Mt. Shasta						0% (0; 12)	9.8% (2; 10)	5.0% (2; 47)	8.8% (1; 16)	0% (0; 18)	0% (0; 22)	2.3% (1; 86)
Solano	Kaiser Foundation Hospital – Rehabilitation Center Vallejo	0% (0; 3)		0% (0; 21)			0% (0; 72)	7.9% (19; 142)	3.6% (21; 412)	9.2% (28; 338)	2.2% (5; 313)	3.3% (7; 147)	3.5% (20; 474)
	North Bay Medical Center							17.0% (10; 34) Worse	1.3% (2; 189)	4.2% (2; 46)	2.1% (2; 113)	2.6% (1; 36)	4.6% (12; 243)
	North Bay VacaValley Hospital							11.8% (6; 22)	4.7% (8; 111)	9.3% (5; 51)	1.7% (1; 79)	10.5% (4; 42) Worse	3.6% (11; 203)
	Sutter Solano Medical Center						0% (0; 5)	9.0% (7; 44)	2.8% (5; 192)	5.3% (3; 81)	1.5% (1; 63)	2.8% (1; 33)	10.7% (13; 140) Worse
Sonoma	Healdsburg District Hospital								10.1% (3; 19)	20.5% (4; 18)	11.8% (1; 6)	0% (0; 10)	9.3% (4; 59)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Kaiser Foundation Hospital – Santa Rosa		0% (0; 7)				0% (0; 51)	3.1% (5; 129) Better	4.2% (11; 177)	6.5% (9; 160)	4.6% (7; 147)	3.0% (4; 112)	4.8% (12; 281)
	Palm Drive Hospital							0% (0; 3)	1.4% (1; 47)	9.9% (2; 18)	0% (0; 42)	0% (0; 26) Better	1.5% (1; 67)
	Petaluma Valley Hospital						0% (0; 8)	2.9% (3; 58)	2.4% (3; 73)	7.3% (5; 65)	0% (0; 63)	3.4% (1; 38)	3.5% (5; 119)
	Santa Rosa Memorial Hospital – Montgomery			0% (0; 10)	10.7% (5; 33)	1.8% (7; 255)	0% (0; 33)	8.2% (22; 259)	3.6% (13; 273)	13.3% (40; 247)	1.9% (4; 182)	5.1% (8; 143)	5.3% (15; 264)
	Sonoma Valley Hospital							3.9% (3; 31)	0% (0; 56)	7.9% (4; 41)	6.1% (3; 38)	0% (0; 38)	2.1% (2; 64)
	Sutter Medical Center of Santa Rosa			0% (0; 15)		2.9% (8; 278)	0% (0; 36)	7.0% (9; 190)	6.3% (6; 133)	14.7% (9; 47)	0.9% (1; 75)	0% (0; 28)	3.0% (2; 83)
Stanislaus	Doctors Medical Center			7.5% (1; 29)	10.6% (3; 35)	1.9% (10; 570)	0.8% (1; 154)	7.4% (26; 423)	3.2% (13; 435)	12.8% (33; 225)	2.9% (6; 218)	4.8% (4; 102)	6.2% (16; 322)
	Emanuel Medical Center, Inc.						0% (0; 8)	6.5% (9; 85)	3.6% (16; 382)	11.5% (10; 147)	1.8% (3; 182)	1.9% (1; 50)	6.9% (32; 475) Worse
	Memorial Hospital Medical Center – Modesto			0% (0; 31)	7.0% (4; 37)	2.1% (17; 886)	0.6% (1; 165)	7.3% (32; 478)	4.1% (29; 539)	11.2% (52; 372)	2.4% (8; 342)	3.9% (13; 257)	5.6% (31; 603)
	Oak Valley District Hospital							4.2% (2; 26)	2.7% (3; 99)	8.4% (2; 35)	2.5% (1; 41)	0% (0; 12)	6.4% (4; 63)
Sutter	Fremont Medical Center								5.4% (6; 79)	12.2% (4; 56)	4.4% (3; 69)	0% (0; 30)	6.3% (9; 216)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
Tehama	Saint Elizabeth Community Hospital							4.1% (1; 13)	2.4% (3; 114)	9.1% (5; 74)	6.6% (4; 65)	4.7% (2; 45)	4.9% (7; 178)
Trinity	Trinity Hospital							0% (0; 6)	6.4% (1; 17)	47.8% (2; 11)	0% (0; 18)		3.8% (1; 49)
Tulare	Kaweah Delta Medical Center			0% (0; 15)	21.8% (1; 6)	1.4% (10; 1,273)	0% (0; 104)	6.7% (15; 360)	2.2% (9; 387)	16.0% (37; 293) Worse	3.4% (7; 228)	5.9% (8; 171)	3.7% (18; 542)
	Sierra View District Hospital					0% (0; 3)	0% (0; 3)	9.2% (9; 69)	5.7% (18; 232) Worse	15.1% (18; 176)	4.1% (6; 156)	5.9% (3; 58)	7.9% (37; 412) Worse
	Tulare District Hospital							21.5% (9; 28) Worse	10.3% (10; 123) Worse	10.8% (8; 75)	6.3% (3; 70)	2.5% (1; 43)	6.4% (7; 174)
Tuolumne	Sonora Regional Medical Center – Greenley			0% (0; 8)			0% (0; 18)	5.4% (4; 34)	3.9% (8; 128)	11.0% (4; 55)	2.7% (3; 105)	2.9% (2; 70)	2.3% (6; 219) Better
Ventura	Community Memorial Hospital – San Buenaventura			0% (0; 19)	0% (0; 25)	1.7% (7; 346)	0% (0; 74)	7.4% (17; 262)	2.8% (10; 316)	9.1% (19; 229)	1.1% (3; 173)	0% (0; 149) Better	3.5% (14; 314)
	Los Robles Hospital and Medical Center			0% (0; 22)	13.2% (5; 50)	1.8% (6; 439)	0% (0; 36)	8.6% (24; 310)	5.7% (20; 320) Worse	12.8% (22; 187)	1.1% (2; 207)	2.4% (3; 148)	2.3% (5; 214)
	Ojai Valley Community Hospital							17.5% (5; 17)	9.2% (1; 13)	9.7% (2; 21)	0% (0; 22)	0% (0; 18)	6.3% (3; 61)
	Saint John’s Pleasant Valley Hospital						0% (0; 3)	2.7% (2; 36)	1.4% (3; 135)	12.4% (10; 65)	0.8% (1; 90)	3.3% (3; 88)	5.0% (9; 139)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2008

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.4% (25; 466)	5% (34; 684)	2.1% (55; 2,619)	7.5% (1,022; 13,616)	1.8% (941; 51,163)	0.6% (49; 8,042)	7.5% (3,462; 46,066)	3.4% (2,928; 86,601)	10.6% (5,322; 50,396)	2.4% (1,161; 48,430)	2.7% (642; 23,627)	4.8% (3,793; 79,667)
	Saint John's Regional Medical Center			0% (0; 21)	11.1% (4; 35)	1.5% (8; 267)	0% (0; 75)	4.5% (11; 223)	1.9% (6; 261)	9.7% (26; 239)	2.9% (9; 214)	0% (0; 71) Better	3.9% (17; 357)
	Simi Valley Hospital and Healthcare Services – Sycamore				78.3% (1; 4)		0% (0; 7)	5.1% (5; 42)	4.8% (9; 147)	10.5% (7; 76)	1.8% (3; 87)	2.6% (1; 54)	2.6% (4; 156)
	Ventura County Medical Center				9.0% (5; 31)			20.0% (6; 27) Worse	3.2% (5; 139)	12.5% (19; 121)	2.7% (2; 92)	2.3% (1; 57)	12.5% (21; 183) Worse
Yolo	Sutter Davis Hospital						0% (0; 9)	9.0% (3; 10)	1.5% (1; 49)	5.4% (3; 43)	1.4% (1; 46)	0% (0; 31)	4.3% (7; 132)
	Woodland Memorial Hospital						0% (0; 6)	10.5% (4; 29)	3.7% (5; 94)	11.2% (5; 86)	1.8% (2; 79)	0% (0; 38)	4.6% (5; 131)
Yuba	Rideout Memorial Hospital	0% (0; 3)		0% (0; 5)		2.1% (5; 402)	0% (0; 60)	6.3% (13; 271)	4.2% (13; 307)	14.6% (14; 126)	1.8% (3; 170)	2.7% (2; 84)	6.2% (16; 301)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2008. See [Technical Note](#) for details about methodology.

**Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms:

Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

Number of Deaths: number of patients that died in this hospital.

Number of Cases: total number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.