

AMI Report: St. Francis Medical Center-Lynwood - Model A

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	146	29	22.0	19.9	15.1	19.8	14.3	25.4	0.062
1992	14.5	156	23	23.9	14.7	15.3	13.9	8.9	18.9	0.470
1993	13.9	114	18	17.1	15.8	15.0	14.7	8.9	20.5	0.441
1994	13.0	188	29	21.7	15.4	11.5	17.4	12.5	22.2	0.053
1995	12.8	184	24	25.9	13.0	14.1	11.9	7.6	16.1	0.382
1996	12.2	209	29	26.8	13.9	12.8	13.2	9.1	17.2	0.347
1997	12.0	182	14	18.9	7.7	10.4	8.9	4.0	13.8	0.130
1998	12.1	145	7	13.9	4.8	9.6	6.1	0.4	11.8	0.022

Model A is a conservative model with risk factors of chronic and personal characters

AMI Report: St. Francis Medical Center-Lynwood - Model B

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	146	29	23.4	19.9	16.0	18.7	13.9	23.4	0.090
1992	14.5	156	23	23.0	14.7	14.7	14.5	9.8	19.2	0.544
1993	13.9	113	18	16.9	15.9	15.0	14.8	9.6	20.0	0.420
1994	12.9	183	29	20.6	15.8	11.3	18.2	13.6	22.8	0.020
1995	12.8	174	23	25.5	13.2	14.7	11.5	7.7	15.3	0.303
1996	12.2	204	29	26.6	14.2	13.0	13.3	9.5	17.0	0.319
1997	12.0	182	14	18.5	7.7	10.2	9.1	4.4	13.7	0.133
1998	12.1	144	7	13.2	4.9	9.2	6.4	0.9	11.9	0.024

Model B is a comprehensive model with more acute risk factors added in