

AMI Report: Riverside Community Hospital - Model A

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	191	31	28.5	16.2	14.9	16.4	11.6	21.2	0.325
1992	14.5	207	27	28.0	13.0	13.5	14.0	9.3	18.6	0.466
1993	13.9	157	15	19.3	9.6	12.3	10.8	5.4	16.3	0.163
1994	13.0	179	24	22.6	13.4	12.6	13.7	9.1	18.4	0.410
1995	12.8	211	32	26.8	15.2	12.7	15.3	11.0	19.5	0.152
1996	12.2	204	21	25.1	10.3	12.3	10.2	6.0	14.3	0.205
1997	12.0	241	29	26.1	12.0	10.8	13.4	9.2	17.5	0.295
1998	12.1	207	32	26.8	15.5	13.0	14.4	10.4	18.4	0.152

Model A is a conservative model with risk factors of chronic and personal characters

AMI Report: Riverside Community Hospital - Model B

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	191	31	27.6	16.2	14.4	17.0	12.5	21.4	0.236
1992	14.5	207	27	27.1	13.0	13.1	14.5	10.2	18.7	0.550
1993	13.9	157	15	20.3	9.6	12.9	10.3	5.4	15.2	0.092
1994	12.9	179	24	27.6	13.4	15.4	11.3	7.5	15.0	0.225
1995	12.8	211	32	28.4	15.2	13.5	14.4	10.7	18.2	0.226
1996	12.2	204	21	25.0	10.3	12.2	10.3	6.4	14.1	0.199
1997	12.0	236	28	23.6	11.9	10.0	14.3	10.2	18.4	0.170
1998	12.1	199	31	28.5	15.6	14.3	13.2	9.6	16.7	0.314

Model B is a comprehensive model with more acute risk factors added in